



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



KCNMB4 siRNA (m): sc-146370

BACKGROUND

MaxiK channels are large conductance voltage and Ca^{2+} -activated potassium channels which are formed by tetramers of MaxiK α subunits, which create pores that are used for smooth muscle tone and neuronal excitability. These MaxiK α subunits have the ability to coassemble with MaxiK β subunits that are structurally related and are able to regulate the function of MaxiK α subunits. KCNMB4 (potassium large conductance calcium-activated channel, subfamily M, β member 4), also known as Slo- β -4 or Maxi K channel subunit β -4, is a 210 amino acid multi-pass membrane protein belonging to the KCNMB family. Predominantly expressed in brain, KCNMB4 is a regulatory subunit of the calcium activated potassium MaxiK α channel. KCNMB4 contributes to MaxiK α channel diversity by modulating calcium sensitivity and gating kinetics of MaxiK α .

REFERENCES

1. Berkovic, S.F. 1997. Epilepsy genes and the genetics of epilepsy syndromes: the promise of new therapies based on genetic knowledge. *Epilepsia* 38: S32-S36.
2. Behrens, R., Nolting, A., Reimann, F., Schwarz, M., Waldschütz, R. and Pongs, O. 2000. hKCNMB3 and hKCNMB4, cloning and characterization of two members of the large-conductance calcium-activated potassium channel β subunit family. *FEBS Lett.* 474: 99-106.
3. Brenner, R., Jegla, T.J., Wickenden, A., Liu, Y. and Aldrich, R.W. 2000. Cloning and functional characterization of novel large conductance calcium-activated potassium channel β subunits, hKCNMB3 and hKCNMB4. *J. Biol. Chem.* 275: 6453-6461.
4. Meera, P., Wallner, M. and Toro, L. 2000. A neuronal β subunit (KCNMB4) makes the large conductance, voltage- and Ca^{2+} -activated K^+ channel resistant to charybdotoxin and iberiotoxin. *Proc. Natl. Acad. Sci. USA* 97: 5562-5567.
5. Jin, P., Weiger, T.M., Wu, Y. and Levitan, I.B. 2002. Phosphorylation-dependent functional coupling of hSlo calcium-dependent potassium channel and its h β 4 subunit. *J. Biol. Chem.* 277: 10014-10020.
6. Jin, P., Weiger, T.M. and Levitan, I.B. 2002. Reciprocal modulation between the α and β 4 subunits of hSlo calcium-dependent potassium channels. *J. Biol. Chem.* 277: 43724-43729.
7. Orio, P., Rojas, P., Ferreira, G. and Latorre, R. 2002. New disguises for an old channel: MaxiK channel β -subunits. *News Physiol. Sci.* 17: 156-161.
8. Henney, N.C., Li, B., Elford, C., Reviriego, P., Campbell, A.K., Wann, K.T. and Evans, B.A. 2009. A large-conductance (BK) potassium channel subtype affects both growth and mineralization of human osteoblasts. *Am. J. Physiol., Cell Physiol.* 297: C1397-C1408.
9. Holtzclaw, J.D., Grimm, P.R. and Sansom, S.C. 2010. Intercalated cell BK- α / β 4 channels modulate sodium and potassium handling during potassium adaptation. *J. Am. Soc. Nephrol.* 21: 634-645.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Kcnmb4 (mouse) mapping to 10 D2.

PRODUCT

KCNMB4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KCNMB4 shRNA Plasmid (m): sc-146370-SH and KCNMB4 shRNA (m) Lentiviral Particles: sc-146370-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KCNMB4 siRNA (m) is recommended for the inhibition of KCNMB4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KCNMB4 gene expression knockdown using RT-PCR Primer: KCNMB4 (m)-PR: sc-146370-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.