



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

KCNV2 siRNA (m): sc-146376

BACKGROUND

Voltage-gated potassium channels play an essential role in controlling cellular excitability in the nervous system. They regulate a variety of properties including membrane potential as well as the frequency and structure of action potentials. KCNV2 (potassium channel, subfamily V, member 2), also known as KV11.1, is a 562 amino acid multi-pass membrane protein that belongs to the potassium channel family, the V subfamily and the Kv8.2/KCNV2 sub-subfamily. KCNV2 forms a heteromultimer with KV2.1, KV3.1 and KIR2.1. Considered a potassium channel subunit, KCNV2 modulates channel activity by shifting the threshold and the half-maximal activation to more negative values. KCNV2 is encoded by a gene located on human chromosome 9p24.2 and mouse chromosome 19 C1. Human chromosome 9 consists of about 145 million bases and encodes nearly 900 genes.

REFERENCES

- Salinas, M., de Weille, J., Guillemare, E., Lazdunski, M. and Hugnot, J.P. 1997. Modes of regulation of shab K⁺ channel activity by the Kv8.1 subunit. *J. Biol. Chem.* 272: 8774-8780.
- Wulfsen, I., Hauber, H.P., Schiemann, D., Bauer, C.K. and Schwarz, J.R. 2000. Expression of mRNA for voltage-dependent and inward-rectifying K channels in GH3/B6 cells and rat pituitary. *J. Neuroendocrinol.* 12: 263-272.
- Yamakura, T., Lewohl, J.M. and Harris, R.A. 2001. Differential effects of general anesthetics on G protein-coupled inwardly rectifying and other potassium channels. *Anesthesiology* 95: 144-153.
- Humphray, S.J., Oliver, K., Hunt, A.R., Plumb, R.W., Loveland, J.E., Howe, K.L., Andrews, T.D., Searle, S., Hunt, S.E., Scott, C.E., Jones, M.C., Ainscough, R., Almeida, J.P., Ambrose, K.D., Ashwell, R.I., et al. 2004. DNA sequence and analysis of human chromosome 9. *Nature* 429: 369-374.
- Czirják, G., Tóth, Z.E. and Enyedi, P. 2007. Characterization of the heteromeric potassium channel formed by kv2.1 and the retinal subunit kv8.2 in *Xenopus* oocytes. *J. Neurophysiol.* 98: 1213-1222.
- Wang, X., Xu, R., Abernathey, G., Taylor, J., Alzghoul, M.B., Hannon, K., Hockerman, G.H. and Pond, A.L. 2008. Kv11.1 channel subunit composition includes MinK and varies developmentally in mouse cardiac muscle. *Dev. Dyn.* 237: 2430-2437.
- Jorge, B.S., Campbell, C.M., Miller, A.R., Rutter, E.D., Gurnett, C.A., Vanoye, C.G., George, A.L. and Kearney, J.A. 2011. Voltage-gated potassium channel KCNV2 (Kv8.2) contributes to epilepsy susceptibility. *Proc. Natl. Acad. Sci. USA* 108: 5443-5448.

CHROMOSOMAL LOCATION

Genetic locus: *Kcnv2* (mouse) mapping to 19 C1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

KCNV2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KCNV2 shRNA Plasmid (m): sc-146376-SH and KCNV2 shRNA (m) Lentiviral Particles: sc-146376-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KCNV2 siRNA (m) is recommended for the inhibition of KCNV2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KCNV2 gene expression knockdown using RT-PCR Primer: KCNV2 (m)-PR: sc-146376-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.