



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Kctd12b siRNA (m): sc-146384

BACKGROUND

The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (Poxvirus and zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C₂H₂-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. Kctd12b (potassium channel tetramerisation domain containing 12b) is a 292 amino acid protein that contains a potassium channel tetramerisation domain and a BTB/POZ domain, which suggests a possible role as a transcriptional regulator. The gene encoding Kctd12b maps to mouse chromosome X F3.

REFERENCES

1. Bardwell, V.J. and Treisman, R. 1994. The POZ domain: a conserved protein-protein interaction motif. *Genes Dev.* 8: 1664-1677.
2. Zollman, S., Godt, D., Prive, G.G., Couderc, J.L. and Laski, F.A. 1994. The BTB domain, found primarily in zinc finger proteins, defines an evolutionarily conserved family that includes several developmentally regulated genes in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 91: 10717-10721.
3. Ahmad, K.F., Engel, C.K. and Prive, G.G. 1998. Crystal structure of the BTB domain from PLZF. *Proc. Natl. Acad. Sci. USA* 95: 12123-12128.
4. Ding, X.F., Luo, C., Ren, K.Q., Zhang, J., Zhou, J.L., Hu, X., Liu, R.S., Wang, Y., Gao, X. and Zhang, J. 2008. Characterization and expression of a human KCTD1 gene containing the BTB domain, which mediates transcriptional repression and homomeric interactions. *DNA Cell Biol.* 27: 257-265.
5. Siggs, O.M. and Beutler, B. 2012. The BTB-ZF transcription factors. *Cell Cycle* 11: 3358-3369.

CHROMOSOMAL LOCATION

Genetic locus: Kctd12b (mouse) mapping to X F3.

PRODUCT

Kctd12b siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Kctd12b shRNA Plasmid (m): sc-146384-SH and Kctd12b shRNA (m) Lentiviral Particles: sc-146384-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Kctd12b siRNA (m) is recommended for the inhibition of Kctd12b expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Kctd12b gene expression knockdown using RT-PCR Primer: Kctd12b (m)-PR: sc-146384-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.