



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Keratin 27 siRNA (m): sc-146408

BACKGROUND

Keratin 27 (K27), also known as KRT27, Cytokeratin-27 (CK-27), Keratin-25C (K25C) or KRT25C, is a 459 amino acid cytoplasmic protein that belongs to the intermediate filament family. Existing as a heterotetramer of two type I and two type II keratins, Keratin 27 interacts with Cytokeratin 6 to form filaments. Keratin 27 is essential for the formation of keratin intermediate filaments in the inner root sheath (irs), and for the proper assembly of type I and type II keratin protein complexes. Keratin 27 is strongly expressed in skin and scalp. In the hair follicle, Keratin 27 is expressed in Henle layer, Huxley layer and in the inner root sheath cuticle of the hair follicle. Keratin 27 expression extends from the bulb region up to the point of differentiation into the three layers. In addition, Keratin 27 is present in the medulla of beard hair. The gene that encodes Keratin 27 consists of approximately 5,727 bases and maps to human chromosome 17q21.2.

REFERENCES

1. Hall, J.M., Friedman, L., Guenther, C., Lee, M.K., Weber, J.L., Black, D.M. and King, M.C. 1992. Closing in on a breast cancer gene on chromosome 17q. *Am. J. Hum. Genet.* 50: 1235-1242.
2. Evans, S.C. and Lozano, G. 1997. The Li-Fraumeni syndrome: an inherited susceptibility to cancer. *Mol. Med. Today* 3: 390-395.
3. Kersemaekers, A.M., Hermans, J., Fleuren, G.J. and van de Vijver, M.J. 1998. Loss of heterozygosity for defined regions on chromosomes 3, 11 and 17 in carcinomas of the uterine cervix. *Br. J. Cancer* 77: 192-200.
4. Piura, B., Rabinovich, A. and Yanai-Inbar, I. 2001. Three primary malignancies related to BRCA mutation successively occurring in a BRCA1 185delAG mutation carrier. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 97: 241-244.
5. Minamoto, T., Buschmann, T., Habelhah, H., Matusevich, E., Tahara, H., Boerresen-Dale, A.L., Harris, C., Sidransky, D. and Ronai, Z. 2001. Distinct pattern of p53 phosphorylation in human tumors. *Oncogene* 20: 3341-3347.
6. Rogers, M.A., Winter, H., Langbein, L., Bleiler, R. and Schweizer, J. 2004. The human type I keratin gene family: characterization of new hair follicle specific members and evaluation of the chromosome 17q21.2 gene domain. *Differentiation* 72: 527-540.
7. Langbein, L., Rogers, M.A., Praetzel-Wunder, S., Helmke, B., Schirmacher, P. and Schweizer, J. 2006. K25 (K25irs1), K26 (K25irs2), K27 (K25irs3), and K28 (K25irs4) represent the type I inner root sheath keratins of the human hair follicle. *J. Invest. Dermatol.* 126: 2377-2386.
8. Yu, Z., Gordon, S.W., Nixon, A.J., Bawden, C.S., Rogers, M.A., Wildermoth, J.E., Maqbool, N.J. and Pearson, A.J. 2009. Expression patterns of keratin intermediate filament and keratin associated protein genes in wool follicles. *Differentiation* 77: 307-316.

CHROMOSOMAL LOCATION

Genetic locus: Krt27 (mouse) mapping to 11 D.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Keratin 27 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Keratin 27 shRNA Plasmid (m): sc-146408-SH and Keratin 27 shRNA (m) Lentiviral Particles: sc-146408-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Keratin 27 siRNA (m) is recommended for the inhibition of Keratin 27 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Keratin 27 gene expression knockdown using RT-PCR Primer: Keratin 27 (m)-PR: sc-146408-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.