



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Keratin 74 siRNA (m): sc-146422

BACKGROUND

The Keratin multigene family is made of "soft" epithelial cytokeratins and "hard" hair Keratins. While the epithelial cytokeratins are involved in the layering and formation of epithelia, the hair Keratins are responsible for creating nails and hair. There are two types of Keratins: the acidic class I Keratin proteins and the basic/neutral class II Keratin proteins. Keratin 74 (KRT74), also known as K6IRS4, KRT5C or KRT6IRS4, is a 529 amino acid protein that is highly expressed in scalp hair follicles. Specifically, Keratin 74 is found in the Huxley layer of the inner root sheath (IRS). Woolly hair autosomal dominant (ADWH) is a rare disorder caused by defects in the Keratin 74 gene which causes fine and tightly curled hair that stops growing after a few inches. Only affecting the scalp, progressive hair loss begins at early adulthood and complete baldness occurs after 30 years. The gene encoding Keratin 74 maps to human chromosome 12q13.13.

REFERENCES

1. Rogers, M.A., Langbein, L., Praetzel, S., Moll, I., Krieg, T., Winter, H. and Schweizer, J. 1997. Sequences and differential expression of three novel human type-II hair keratins. *Differentiation* 61: 187-194.
2. Rogers, M.A., Winter, H., Langbein, L., Wolf, C. and Schweizer, J. 2000. Characterization of a 300 kbp region of human DNA containing the type II hair keratin gene domain. *J. Invest. Dermatol.* 114: 464-472.
3. Langbein, L., Rogers, M.A., Winter, H., Praetzel, S. and Schweizer, J. 2001. The catalog of human hair keratins. II. Expression of the six type II members in the hair follicle and the combined catalog of human type I and II keratins. *J. Biol. Chem.* 276: 35123-35132.
4. Hesse, M., Magin, T.M. and Weber, K. 2001. Genes for intermediate filament proteins and the draft sequence of the human genome: novel keratin genes and a surprisingly high number of pseudogenes related to keratin genes 8 and 18. *J. Cell Sci.* 114: 2569-2575.
5. Langbein, L. and Schweizer, J. 2005. Keratins of the human hair follicle. *Int. Rev. Cytol.* 243: 1-78.
6. Rogers, M.A., Edler, L., Winter, H., Langbein, L., Beckmann, I. and Schweizer, J. 2005. Characterization of new members of the human type II keratin gene family and a general evaluation of the keratin gene domain on chromosome 12q13.13. *J. Invest. Dermatol.* 124: 536-544.
7. Langbein, L., Rogers, M.A., Praetzel-Wunder, S., Helmke, B., Schirmacher, P. and Schweizer, J. 2006. K25 (K25irs1), K26 (K25irs2), K27 (K25irs3), and K28 (K25irs4) represent the type I inner root sheath keratins of the human hair follicle. *J. Invest. Dermatol.* 126: 2377-2386.
8. Shimomura, Y., Wajid, M., Petukhova, L., Kurban, M. and Christiano, A.M. 2010. Autosomal-dominant woolly hair resulting from disruption of Keratin 74 (KRT74), a potential determinant of human hair texture. *Am. J. Hum. Genet.* 86: 632-638.
9. Wasif, N., Naqvi, S.K., Basit, S., Ali, N., Ansar, M. and Ahmad, W. 2011. Novel mutations in the Keratin 74 (KRT74) gene underlie autosomal dominant woolly hair/hypotrichosis in Pakistani families. *Hum. Genet.* 129: 419-424.

CHROMOSOMAL LOCATION

Genetic locus: Krt74 (mouse) mapping to 15 F2.

PRODUCT

Keratin 74 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Keratin 74 shRNA Plasmid (m): sc-146422-SH and Keratin 74 shRNA (m) Lentiviral Particles: sc-146422-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Keratin 74 siRNA (m) is recommended for the inhibition of Keratin 74 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Keratin 74 gene expression knockdown using RT-PCR Primer: Keratin 74 (m)-PR: sc-146422-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.