



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CEP162 siRNA (m): sc-146449

BACKGROUND

CEP162 (Centrosomal protein of 162 kDa), also known as KIAA1009 or QN1 (quail neuroretina 1), is a 1,403 amino acid protein in the CEP162 family and containing four coiled coil domains. CEP162 is an axoneme-associated protein required to promote the assembly of the transition zone at the base of primary cilia. CEP162 is localized at the distal ends of the centrioles before ciliogenesis and acts by specifically recognizing and binding the axonemal microtubule, effectively restricting zone formation specifically at the cilia base. During mitosis, CEP162 is located at the spindle poles and the centrosomes and is essential for chromosome segregation and spindle assembly, interacting with α -Tubulin. CEP162 interacts with CEP290 and is required to mediate its association with microtubules. The CEP162 gene is widely conserved, including mouse, rat, zebrafish, canine and bovine.

REFERENCES

1. Munnia, A., Schütz, N., Romeike, B.F., Maldener, E., Glass, B., Maas, R., Nastainczyk, W., Feiden, W., Fischer, U. and Meese, E. 2001. Expression, cellular distribution and protein binding of the glioma amplified sequence (GAS41), a highly conserved putative transcription factor. *Oncogene* 20: 4853-4863.
2. Jia, L., Young, M.F., Powell, J., Yang, L., Ho, N.C., Hotchkiss, R., Robey, P.G. and Francomano, C.A. 2002. Gene expression profile of human bone marrow stromal cells: high-throughput expressed sequence tag sequencing analysis. *Genomics* 79: 7-17.
3. Tomsig, J.L., Snyder, S.L. and Creutz, C.E. 2003. Identification of targets for calcium signaling through the copine family of proteins. Characterization of a coiled-coil copine-binding motif. *J. Biol. Chem.* 278: 10048-10054.
4. Leon, A., Omri, B., Gely, A., Klein, C. and Crisanti, P. 2006. QN1/KIAA1009: a new essential protein for chromosome segregation and mitotic spindle assembly. *Oncogene* 25: 1887-1895.
5. Jakobsen, L., Vanselow, K., Skogs, M., Toyoda, Y., Lundberg, E., Poser, I., Falkenby, L.G., Bennetzen, M., Westendorf, J., Nigg, E.A., Uhlen, M., Hyman, A.A. and Andersen, J.S. 2011. Novel asymmetrically localizing components of human centrosomes identified by complementary proteomics methods. *EMBO J.* 30: 1520-1535.
6. Wang, W.J., Tay, H.G., Soni, R., Perumal, G.S., Goll, M.G., Macaluso, F.P., Asara, J.M., Amack, J.D. and Tsou, M.F. 2013. CEP162 is an axoneme-recognition protein promoting ciliary transition zone assembly at the cilia base. *Nat. Cell Biol.* 15: 591-601.
7. Awata, T., Yamashita, H., Kurihara, S., Morita-Ohkubo, T., Miyashita, Y., Katayama, S., Mori, K., Yoneya, S., Kohda, M., Okazaki, Y., Maruyama, T., Shimada, A., Yasuda, K., Nishida, N., Tokunaga, K. and Koike, A. 2014. A genome-wide association study for diabetic retinopathy in a Japanese population: potential association with a long intergenic non-coding RNA. *PLoS ONE* 9: e111715.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Cep162 (mouse) mapping to 9 E3.1.

PRODUCT

CEP162 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CEP162 shRNA Plasmid (m): sc-146449-SH and CEP162 shRNA (m) Lentiviral Particles: sc-146449-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CEP162 siRNA (m) is recommended for the inhibition of CEP162 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CEP162 gene expression knockdown using RT-PCR Primer: CEP162 (m)-PR: sc-146449-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.