



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

KY peptidase siRNA (m): sc-146614

BACKGROUND

Filamins are Actin-binding proteins which contain an N-terminal Actin-binding domain, a membrane glycoprotein domain and a C-terminal self-association domain. Filamins help reshape the cytoskeleton by forming flexible cross-links between two Actin filaments, which maintain membrane integrity during force application. Filamin 2, also designated Filamin C, is a skeletal- and cardiac-muscle specific form of Filamin, which binds γ -sarcoglycan and δ -sarcoglycan, but not α -sarcoglycan or β -sarcoglycan. KY peptidase (kyphoscoliosis peptidase) is a 561 amino acid cytoskeleton protease that interacts with several sarcomeric cytoskeletal proteins, including Filamin 2. KY peptidase probably plays a role in the maturation, function and stabilization of the neuromuscular junction. KY-null mouse mutants exhibit distinct irregular subcellular Filamin 2 localization, suggesting that KY peptidase deficiency may be the cause of several types of limb-girdle muscular dystrophies.

REFERENCES

1. Blanco, G., Nikitopoulou, A., Kraus, M., Mason, R.M., Coulton, G.R. and Brown, S.D. 1998. A STS content physical and transcription map across the KY, kyphoscoliosis, nonrecombinant region. *Genomics* 54: 415-423.
2. van der Ven, P.F., Obermann, W.M., Lemke, B., Gautel, M., Weber, K. and Fürst, D.O. 2000. Characterization of muscle filamin isoforms suggests a possible role of γ -Filamin/ABP-L in sarcomeric Z-disc formation. *Cell Motil. Cytoskeleton* 45: 149-162.
3. Thompson, T.G., Chan, Y.M., Hack, A.A., Brosius, M., Rajala, M., Lidov, H.G., McNally, E.M., Watkins, S. and Kunkel, L.M. 2000. Filamin 2 (FLN2): a muscle-specific sarcoglycan interacting protein. *J. Cell Biol.* 148: 115-126.
4. Blanco, G., Coulton, G.R., Biggin, A., Grainge, C., Moss, J., Barrett, M., Berquin, A., Marechal, G., Skynner, M., van Mier, P., Nikitopoulou, A., Kraus, M., Ponting, C.P., Mason, R.M. and Brown, S.D. 2001. The kyphoscoliosis (KY) mouse is deficient in hypertrophic responses and is caused by a mutation in a novel muscle-specific protein. *Hum. Mol. Genet.* 10: 9-16.
5. Murray, J.T., Campbell, D.G., Pegg, M., Mora, A., Alfonso, M. and Cohen, P. 2004. Identification of Filamin C as a new physiological substrate of PKB α using KESTREL. *Biochem. J.* 384: 489-494.
6. Beatham, J., Romero, R., Townsend, S.K., Hacker, T., van der Ven, P.F. and Blanco, G. 2004. Filamin C interacts with the muscular dystrophy KY protein and is abnormally distributed in mouse KY deficient muscle fibres. *Hum. Mol. Genet.* 13: 2863-2874.
7. Anastasi, G., Cutroneo, G., Trimarchi, F., Santoro, G., Bruschetta, D., Bramanti, P., Pisani, A. and Favaloro, A. 2004. Evaluation of sarcoglycans, vinculin-talin-integrin system and Filamin 2 in α - and γ -sarcoglycanopathy: an immunohistochemical study. *Int. J. Mol. Med.* 14: 989-999.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 605739. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Ky (mouse) mapping to 9 F1.

PRODUCT

KY peptidase siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KY peptidase shRNA Plasmid (m): sc-146614-SH and KY peptidase shRNA (m) Lentiviral Particles: sc-146614-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KY peptidase siRNA (m) is recommended for the inhibition of KY peptidase expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KY peptidase gene expression knockdown using RT-PCR Primer: KY peptidase (m)-PR: sc-146614-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.