



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LAT2 siRNA (m): sc-146657

BACKGROUND

L-amino acid transporter protein-2 (LAT2), a non-glycosylated membrane protein, complexes with CD98 to contribute to reabsorption of neutral amino acids in renal epithelia and blood-tissue barriers. The gene encoding LAT2 is expressed primarily in the kidney, but also to a lesser extent in placenta, brain, liver, spleen, skeletal muscle, heart, small intestine, and lung. Transfection with the antisense sequence of LAT2 suggests that LAT2 expression plays a major role in net basolateral efflux of cysteine, and points to LAT2 as a candidate gene to modulate cysteine reabsorption. In addition, the CD98/LAT2 heterodimer associates with Integrin β 1 in intestinal epithelial cells, where ligand binding to CD98 and another cell surface molecule, ICAM-1 differentially regulates LAT2 activity, suggesting a novel mechanism by which events like cell adhesion may affect amino acid transport activity.

REFERENCES

1. Pineda, M., et al. 1999. Identification of a membrane protein, LAT2, that Co-expresses with 4F2 heavy chain, an L-type amino acid transport activity with broad specificity for small and large zwitterionic amino acids. *J. Biol. Chem.* 274: 19738-19744.
2. Segawa, H., et al. 1999. Identification and functional characterization of a Na⁺-independent neutral amino acid transporter with broad substrate selectivity. *J. Biol. Chem.* 274: 19745-19751.
3. Fernandez, E., et al. 2003. Basolateral LAT2 has a major role in the transepithelial flux of L-cystine in the renal proximal tubule cell line OK. *J. Am. Soc. Nephrol.* 14: 837-847.
4. Liu, X., et al. 2003. CD98 and intracellular adhesion molecule 1 regulate the activity of amino acid transporter LAT2 in polarized intestinal epithelia. *J. Biol. Chem.* 278: 23672-23677.
5. Pinho, M.J., et al. 2003. Organ-specific overexpression of renal LAT2 and enhanced tubular L-DOPA uptake precede the onset of hypertension. *Hypertension* 42: 613-618.
5. Kim do, K., et al. 2004. System L-amino acid transporters are differently expressed in rat astrocyte and C6 glioma cells. *Neurosci. Res.* 50: 437-446.
6. Soares-Da-Silva, P., et al. 2004. Cloning and gene silencing of LAT2, the L-3,4-dihydroxyphenylalanine (L-DOPA) transporter, in pig renal LLC-PK1 epithelial cells. *FASEB J.* 18: 1489-1498.
7. Shennan, D.B., et al. 2004. L-leucine transport in human breast cancer cells (MCF-7 and MDA-MB-231): kinetics, regulation by estrogen and molecular identity of the transporter. *Biochim. Biophys. Acta* 1664: 206-216.
8. Pinho, M.J., et al. 2004. Over-expression of renal LAT1 and LAT2 and enhanced L-DOPA uptake in SHR immortalized renal proximal tubular cells. *Kidney Int.* 66: 216-226.

CHROMOSOMAL LOCATION

Genetic locus: Slc7a8 (mouse) mapping to 14 C3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

LAT2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LAT2 shRNA Plasmid (m): sc-146657-SH and LAT2 shRNA (m) Lentiviral Particles: sc-146657-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LAT2 siRNA (m) is recommended for the inhibition of LAT2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LAT2 gene expression knockdown using RT-PCR Primer: LAT2 (m)-PR: sc-146657-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.