



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LDH-AL6B siRNA (m): sc-146690

BACKGROUND

The lactate dehydrogenase family (LDH) consists of three members, designated LDH-A, LDH-B and LDH-C, all of which work in concert to catalyze the final step of anaerobic glycolysis, namely the conversion of L-lactate and NAD⁺ to pyruvate and NADH. Each family member displays a specific tissue distribution pattern, with LDH-A present in muscle and LDH-B present in heart, while LDH-C expression is confined to testis and sperm. LDH-AL6B (lactate dehydrogenase A-like 6B), also known as LDHL or LDHAL6, is a 381 amino acid testis-specific protein that functions in a similar manner to LDH-A, specifically catalyzing the NAD⁺-dependent formation of pyruvate and NADH.

REFERENCES

1. Edwards, Y.H., et al. 1987. Locus determining the human sperm-specific lactate dehydrogenase, LDHC, is syntenic with LDHA. *Dev. Genet.* 8: 219-232.
2. LeVan, K.M., et al. 1991. Properties of human testis-specific lactate dehydrogenase expressed from *Escherichia coli*. *Biochem. J.* 273: 587-592.
3. Kanno, T., et al. 1995. Lactate dehydrogenase M-subunit deficiencies: clinical features, metabolic background, and genetic heterogeneities. *Muscle Nerve Suppl.* 3: 54-60.
4. Kopperschlager, G., et al. 1996. Methods for the separation of lactate dehydrogenases and clinical significance of the enzyme. *J. Chromatogr. B, Biomed. Appl.* 684: 25-49.
5. Auerbach, G., et al. 1998. Lactate dehydrogenase from the hyperthermophilic bacterium *Thermotoga maritima*: the crystal structure at 2.1 Å resolution reveals strategies for intrinsic protein stabilization. *Structure* 6: 769-781.
6. Niwakawa, M., et al. 2001. The role of tumor markers in the treatment of germ cell tumor. *Gan To Kagaku Ryoho* 28: 1159-1165.
7. Pioli, P.A., et al. 2002. Lactate dehydrogenase is an AU-rich element-binding protein that directly interacts with AUF1. *J. Biol. Chem.* 277: 35738-35745.
8. Chen, X., et al. 2009. Identification of a novel human lactate dehydrogenase gene LDH-AL6A, which activates transcriptional activities of AP1(PMA). *Mol. Biol. Rep.* 36: 669-76.

CHROMOSOMAL LOCATION

Genetic locus: Ldhal6b (mouse) mapping to 17 A1.

PRODUCT

LDH-AL6B siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LDH-AL6B shRNA Plasmid (m): sc-146690-SH and LDH-AL6B shRNA (m) Lentiviral Particles: sc-146690-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LDH-AL6B siRNA (m) is recommended for the inhibition of LDH-AL6B expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LDH-AL6B gene expression knockdown using RT-PCR Primer: LDH-AL6B (m)-PR: sc-146690-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.