



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Lipocalin-6 siRNA (m): sc-146751

BACKGROUND

The Lipocalin family is composed of structurally conserved hydrophobic ligand and binding proteins and is represented in all major taxonomic groups from prokaryotes to primates. Members of the lipocalin family are characterized by several common molecular-recognition properties: the ability to bind a range of small hydrophobic molecules, binding to specific cell-surface receptors and the formation of complexes with soluble macromolecules. Lipocalin-6, also known as LCN5, hLcn5, UNQ643 or LCN6, is a 163 amino acid protein that is predominantly expressed in epididymis. Lipocalin-6 localizes to the head and tail of spermatozoa, with the highest concentration on the post-acrosomal region of the head. Belonging to the calycin superfamily, Lipocalin-6 may play a role in sperm maturation. The gene encoding Lipocalin-6 maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome.

REFERENCES

- Schmale, H., Holtgreve-Grez, H. and Christiansen, H. 1990. Possible role for salivary gland protein in taste reception indicated by homology to lipophilic-ligand carrier proteins. *Nature* 343: 366-369.
- Flower, D.R. 1995. Multiple molecular recognition properties of the lipocalin protein family. *J. Mol. Recognit.* 8: 185-195.
- Flower, D.R. 1996. The lipocalin protein family: structure and function. *Biochem. J.* 318: 1-14.
- Yenugu, S., Hamil, K.G., Birse, C.E., Ruben, S.M., French, F.S. and Hall, S.H. 2003. Antibacterial properties of the sperm-binding proteins and peptides of human epididymis 2 (HE2) family; salt sensitivity, structural dependence and their interaction with outer and cytoplasmic membranes of *Escherichia coli*. *Biochem. J.* 372: 473-483.
- Hamil, K.G., Liu, Q., Sivashanmugam, P., Anbalagan, M., Yenugu, S., Soundararajan, R., Grossman, G., Rao, A.J., Birse, C.E., Ruben, S.M., Richardson, R.T., Zhang, Y.L., O'Rand, M.G., Petrusz, P., French, F.S., et al. 2003. LCN6, a novel human epididymal lipocalin. *Reprod. Biol. Endocrinol.* 1: 112.
- Suzuki, K., Lareyre, J.J., Sánchez, D., Gutierrez, G., Araki, Y., Matusik, R.J. and Orgebin-Crist, M.C. 2004. Molecular evolution of epididymal lipocalin genes localized on mouse chromosome 2. *Gene* 339: 49-59.
- Grzyb, J., Latowski, D. and Strzałka, K. 2006. Lipocalins-a family portrait. *J. Plant Physiol.* 163: 895-915.
- Suzuki, K., Yu, X., Chaurand, P., Araki, Y., Lareyre, J.J., Caprioli, R.M., Orgebin-Crist, M.C. and Matusik, R.J. 2007. Epididymis-specific lipocalin promoters. *Asian J. Androl.* 9: 515-521.
- Guo, C., Lian, Y., Liu, Q., Liu, J., Zhang, Y. and Lin, D. 2010. Soluble expression and characterization of a mouse epididymis-specific protein Lipocalin-6. *Protein Expr. Purif.* 69: 64-67.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Lcn6 (mouse) mapping to 2 A3.

PRODUCT

Lipocalin-6 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lipocalin-6 shRNA Plasmid (m): sc-146751-SH and Lipocalin-6 shRNA (m) Lentiviral Particles: sc-146751-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Lipocalin-6 siRNA (m) is recommended for the inhibition of Lipocalin-6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Lipocalin-6 gene expression knockdown using RT-PCR Primer: Lipocalin-6 (m)-PR: sc-146751-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.