



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



Liprin α 4 siRNA (m): sc-146755

BACKGROUND

Liprins interact with members of the leukocyte common antigen-related (LAR) family of transmembrane protein tyrosine phosphatases, which are implicated in axon guidance and mammary gland development. Liprins are multivalent proteins that form complex structures and act as scaffolds for the recruitment and anchoring of LAR phosphatases. Based on sequence similarities and binding characteristics, liprins are subdivided into α -type and β -type liprins. Both α - and β -liprins homodimerize via their N-terminal, coiled coil regions. Liprin α 4, also known as PTPRF-interacting protein α 4, regulates the disassembly of focal adhesions. Liprin α 4 is expressed in heart, brain and skeletal muscle. Due to alternative splicing, a second isoform exists for Liprin α 4 that lacks amino acids 444-452.

REFERENCES

1. Serra-Pagès, C., Medley, Q.G., Tang, M., Hart, A. and Streuli, M. 1998. Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting proteins. *J. Biol. Chem.* 273: 15611-15620.
2. Zhen, M. and Jin, Y. 1999. The liprin protein SYD-2 regulates the differentiation of presynaptic termini in *C. elegans*. *Nature* 401: 371-375.
3. Kaufmann, N., DeProto, J., Ranjan, R., Wan, H. and Van Vactor, D. 2002. *Drosophila* liprin- α and the receptor phosphatase Dlar control synapse morphogenesis. *Neuron* 34: 27-38.
4. Ko, J., Kim, S., Valtschanoff, J.G., Shin, H., Lee, J.R., Sheng, M., Premont, R.T., Weinberg, R.J. and Kim, E. 2003. Interaction between liprin- α and GlT1 is required for AMPA receptor targeting. *J. Neurosci.* 23: 1667-1677.
5. Ko, J., Na, M., Kim, S., Lee, J.R. and Kim, E. 2003. Interaction of the ERC family of RIM-binding proteins with the liprin- α family of multidomain proteins. *J. Biol. Chem.* 278: 42377-42385.
6. Katoh, M. and Katoh, M. 2003. Identification and characterization of human PPFIA4 gene in silico. *Int. J. Mol. Med.* 12: 1009-1014.
7. Miller, K.E., DeProto, J., Kaufmann, N., Patel, B.N., Duckworth, A. and Van Vactor, D. 2005. Direct observation demonstrates that liprin- α is required for trafficking of synaptic vesicles. *Curr. Biol.* 15: 684-689.
8. Hofmeyer, K., Maurel-Zaffran, C., Sink, H. and Treisman, J.E. 2006. Liprin- α has LAR-independent functions in R7 photoreceptor axon targeting. *Proc. Natl. Acad. Sci. USA* 103: 11595-11600.
9. Choe, K.M., Prakash, S., Bright, A. and Clandinin, T.R. 2006. Liprin- α is required for photoreceptor target selection in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 103: 11601-11606.

CHROMOSOMAL LOCATION

Genetic locus: Ppfia4 (mouse) mapping to 1 E4.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Liprin α 4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Liprin α 4 shRNA Plasmid (m): sc-146755-SH and Liprin α 4 shRNA (m) Lentiviral Particles: sc-146755-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Liprin α 4 siRNA (m) is recommended for the inhibition of Liprin α 4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Liprin α 4 gene expression knockdown using RT-PCR Primer: Liprin α 4 (m)-PR: sc-146755-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.