



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



LLGL4 siRNA (m): sc-146760

BACKGROUND

In *Drosophila*, Scribble (Scrib), Discs large (Dlg), and Lethal giant larvae (Lgl) function together to regulate epithelial polarity and, significantly, the human homologs of these proteins have been linked to the progression of cancer. LLGL4 (Lethal(2) giant larvae protein homolog 4), also designated Syntaxin-binding protein 5-like or Tomosyn-2, is a 1186 amino acid protein that is expressed in brain hippocampus, kidney, germ cell tumors and lung carcinoma. As a member of the WD repeat L(2)GL family, it is likely that LLGL4 plays a role in vesicle trafficking and exocytosis. LLGL4 contains one v-SNARE coiled-coil homology domain and six WD repeats, which are sites for protein-protein interactions. LLGL4 shares 68% sequence similarity with human LLGL3, also known as Tomosyn, which is an important component in the neurotransmitter release process. There are two isoforms of LLGL4 that are produced as a result of alternative splicing events.

REFERENCES

1. Fujita, Y., Shirataki, H., Sakisaka, T., Asakura, T., Ohya, T., Kotani, H., Yokoyama, S., Nishioka, H., Matsuura, Y., Mizoguchi, A., Scheller, R.H. and Takai, Y. 1998. Tomosyn: a syntaxin-1-binding protein that forms a novel complex in the neurotransmitter release process. *Neuron* 20: 905-915.
2. Yokoyama, S., Shirataki, H., Sakisaka, T. and Takai, Y. 1999. Three splicing variants of tomosyn and identification of their syntaxin-binding region. *Biochem. Biophys. Res. Commun.* 256: 218-222.
3. Humbert, P., Russell, S. and Richardson, H. 2003. Dlg, Scribble and Lgl in cell polarity, cell proliferation and cancer. *Bioessays* 25: 542-553.
4. Katoh, M. and Katoh, M. 2004. Identification and characterization of human LLGL4 gene and mouse Lgl4 gene in silico. *Int. J. Oncol.* 24: 737-742.
5. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 604586. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Groffen, A.J., Jacobsen, L., Schut, D. and Verhage, M. 2005. Two distinct genes drive expression of seven tomosyn isoforms in the mammalian brain, sharing a conserved structure with a unique variable domain. *J. Neurochem.* 92: 554-568.
7. Humbert, P.O., Grzeschik, N.A., Brumby, A.M., Galea, R., Elsum, I. and Richardson, H.E. 2008. Control of tumorigenesis by the Scribble/Dlg/Lgl polarity module. *Oncogene* 27: 6888-6907.

CHROMOSOMAL LOCATION

Genetic locus: Stxbp5l (mouse) mapping to 16 B3.

PRODUCT

LLGL4 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LLGL4 shRNA Plasmid (m): sc-146760-SH and LLGL4 shRNA (m) Lentiviral Particles: sc-146760-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LLGL4 siRNA (m) is recommended for the inhibition of LLGL4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LLGL4 gene expression knockdown using RT-PCR Primer: LLGL4 (m)-PR: sc-146760-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.