



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LTBP-3 siRNA (m): sc-149138

BACKGROUND

LTBP-3 (latent TGF- β -binding protein 3) is a structural component of connective tissue microfibrils that plays an essential role in TGF- β 1 secretion and targeting. LTBP-3 is present at high levels in human heart, skeletal muscle, prostate and ovaries and in certain osteosarcoma and fibroblastic cell lines. Human LTBP-3 localizes to chromosome position 11q12.

REFERENCES

1. Yin, W., Smiley, E., Germiller, J., Mecham, R.P., Florer, J.B., Wenstrup, R.J. and Bonadio, J. 1995. Isolation of a novel latent transforming growth factor- β binding protein gene (LTBP-3). *J. Biol. Chem.* 270: 10147-10160.
2. Yin, W., Smiley, E. and Bonadio, J. 1998. Alternative splicing of LTBP-3. *Biochem. Biophys. Res. Commun.* 245: 454-458.
3. Dabovic, B., Chen, Y., Colarossi, C., Zambuto, L., Obata, H. and Rifkin, D.B. 2002. Bone defects in latent TGF- β binding protein (Ltbp)-3 null mice; a role for Ltbp in TGF- β presentation. *J. Endocrinol.* 175: 129-141.
4. Penttinen, C., Saharinen, J., Weikkolainen, K., Hyytiäinen, M. and Keski-Oja, J. 2002. Secretion of human latent TGF- β -binding protein-3 (LTBP-3) is dependent on co-expression of TGF- β . *J. Cell Sci.* 115: 3457-3468.
5. Chen, Y., Dabovic, B., Annes, J.P. and Rifkin, D.B. 2002. Latent TGF- β binding protein-3 (LTBP-3) requires binding to TGF- β for secretion. *FEBS Lett.* 517: 277-280.
6. Chen, Y., Dabovic, B., Colarossi, C., Santori, F.R., Lilic, M., Vukmanovic, S. and Rifkin, D.B. 2003. Growth retardation as well as spleen and thymus involution in latent TGF- β binding protein (Ltbp)-3 null mice. *J. Cell. Physiol.* 196: 319-325.
7. Kantola, A.K., Keski-Oja, J. and Koli, K. 2005. Induction of human LTBP-3 promoter activity by TGF- β 1 is mediated by Smad3/4 and AP-1 binding elements. *Gene* 363: 142-150.
8. Colarossi, C., Chen, Y., Obata, H., Jurukovski, V., Fontana, L., Dabovic, B. and Rifkin, D.B. 2005. Lung alveolar septation defects in Ltbp-3-null mice. *Am. J. Pathol.* 167: 419-428.

CHROMOSOMAL LOCATION

Genetic locus: Ltbp3 (mouse) mapping to 19 A.

PRODUCT

LTBP-3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LTBP-3 shRNA Plasmid (m): sc-149138-SH and LTBP-3 shRNA (m) Lentiviral Particles: sc-149138-V as alternate gene silencing products.

For independent verification of LTBP-3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149138A, sc-149138B and sc-149138C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LTBP-3 siRNA (m) is recommended for the inhibition of LTBP-3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

LTBP-3 (H-11): sc-390913 is recommended as a control antibody for monitoring of LTBP-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LTBP-3 gene expression knockdown using RT-PCR Primer: LTBP-3 (m)-PR: sc-149138-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.