



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



LYPLA1 siRNA (m): sc-149180

BACKGROUND

The multifunctional lysophospholipids (detergent-like intermediates) found in various biological membranes are regulated by the enzymatic activity of lysophospholipases. Increased levels of lysophospholipids are associated with a host of diseases, including hyperlipidemia and atherosclerosis. LYPLA1 (lysophospholipase 1), also known as APT1 or LPL1, and LYPLA2 (lysophospholipase 2) are lysophospholipases that localize to the cytoplasm and belong to the AB hydrolase 2 family. Both LYPLA1 and LYPLA2 function to enzymatically hydrolyze fatty acids from S-acetylated cysteine residues on a variety of monomeric and micellar substrates, such as H-Ras. Due to their ability to catalytically regulate the overall concentration of lysophospholipids in cellular membranes, LYPLA1 and LYPLA2 may play a crucial role in the development of lysophospholipid-associated disorders. LYPLA1 and LYPLA2 exist as multiple alternatively spliced isoforms that are expressed in tissues throughout the body.

REFERENCES

1. Bohn, E., et al. 1992. Annexin II inhibits calcium-dependent phospholipase A₁ and lysophospholipase but not triacyl glycerol lipase activities of rat liver hepatic lipase. *FEBS Lett.* 296: 237-240.
2. Wang, A., et al. 1999. A specific human lysophospholipase: cDNA cloning, tissue distribution and kinetic characterization. *Biochim. Biophys. Acta* 1437: 157-169.
3. Wang, A. and Dennis, E.A. 1999. Mammalian lysophospholipases. *Biochim. Biophys. Acta* 1439: 1-16.
4. Wang, A., et al. 2000. Subcellular localization and PKC-dependent regulation of the human lysophospholipase A/acyl-protein thioesterase in WISH cells. *Biochim. Biophys. Acta* 1484: 207-214.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605599. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Morello, F., et al. 2004. Differential gene expression of blood-derived cell lines in familial combined hyperlipidemia. *Arterioscler. Thromb. Vasc. Biol.* 24: 2149-2154.

CHROMOSOMAL LOCATION

Genetic locus: Lypla1 (mouse) mapping to 1 A1.

PRODUCT

LYPLA1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LYPLA1 shRNA Plasmid (m): sc-149180-SH and LYPLA1 shRNA (m) Lentiviral Particles: sc-149180-V as alternate gene silencing products.

For independent verification of LYPLA1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149180A, sc-149180B and sc-149180C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LYPLA1 siRNA (m) is recommended for the inhibition of LYPLA1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LYPLA1 gene expression knockdown using RT-PCR Primer: LYPLA1 (m)-PR: sc-149180-PR (20 μ l, 410 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.