



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MAGE-B18 siRNA (m): sc-149219

BACKGROUND

The melanoma-associated antigen (MAGE) family consists of a number of antigens recognized by cytotoxic T lymphocytes. The MAGE genes were initially isolated from different kinds of tumors and, based on their virtually exclusive tumor-specific expression in adult tissues, they have been used as targets for cancer immunotherapy. MAGE genes encode for tumor-rejection antigens that are expressed in tumors of different histologic types and in normal testis and placenta. MAGE-B18 (melanoma-associated antigen B18) is a 343 amino acid protein that contains one MAGE domain and may be involved in tumorigenesis. The gene encoding MAGE-B18 maps to human chromosome X, which contains nearly 153 million base pairs and houses over 1,000 genes.

REFERENCES

1. Graf, L.H. and Ferrone, S. 1989. Human melanoma-associated antigens. Immunol. Ser. 43: 643-679.
2. Merimsky, O., Shoenfeld, Y., Chaitchik, S., Yecheskel, G. and Fishman, P. 1994. Antigens and antibodies in malignant melanoma. Tumour Biol. 15: 188-202.
3. Rogner, U.C., Wilke, K., Steck, E., Korn, B. and Poustka, A. 1995. The melanoma antigen gene (MAGE) family is clustered in the chromosomal band Xq28. Genomics 29: 725-731.
4. Sarantou, T., Chi, D.D., Garrison, D.A., Conrad, A.J., Schmid, P., Morton, D.L. and Hoon, D.S. 1997. Melanoma-associated antigens as messenger RNA detection markers for melanoma. Cancer Res. 57: 1371-1376.
5. Serrano, A., Lethé, B., Delroisse, J.M., Lurquin, C., De Plaen, E., Brasseur, F., Rimoldi, D. and Boon, T. 1999. Quantitative evaluation of the expression of MAGE genes in tumors by limiting dilution of cDNA libraries. Int. J. Cancer 83: 664-669.
6. van Dinten, L.C., Pul, N., van Nieuwpoort, A.F., Out, C.J., Jager, M.J. and van den Elsen, P.J. 2005. Uveal and cutaneous melanoma: shared expression characteristics of melanoma-associated antigens. Invest. Ophthalmol. Vis. Sci. 46: 24-30.
7. Kasuga, C., Nakahara, Y., Ueda, S., Hawkins, C., Taylor, M.D., Smith, C.A. and Rutka, J.T. 2008. Expression of MAGE and GAGE genes in medulloblastoma and modulation of resistance to chemotherapy. Laboratory investigation. J. Neurosurg. Pediatr. 1: 305-313.
8. Tanaka, F., Haraguchi, N., Isikawa, K., Inoue, H. and Mori, M. 2008. Potential role of dendritic cell vaccination with MAGE peptides in gastrointestinal carcinomas. Oncol. Rep. 20: 1111-1116.

CHROMOSOMAL LOCATION

Genetic locus: Mageb18 (mouse) mapping to X C2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

MAGE-B18 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MAGE-B18 shRNA Plasmid (m): sc-149219-SH and MAGE-B18 shRNA (m) Lentiviral Particles: sc-149219-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MAGE-B18 siRNA (m) is recommended for the inhibition of MAGE-B18 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MAGE-B18 gene expression knockdown using RT-PCR Primer: MAGE-B18 (m)-PR: sc-149219-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.