



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Matriptase-2 siRNA (m): sc-149299

## BACKGROUND

Matriptase (also known as MT-SP1, ST14, prostamin and epithin) is a tumor-associated type II transmembrane serine protease that is highly expressed in many human cancer-derived cell lines and is implicated in extracellular matrix remodeling, tumor growth and metastasis. Matriptase performs pleiotropic functions in the development of the epidermis, hair follicles and cellular immune system. Sphingosine 1-phosphate (S1P, SPP), present in serum-derived lipoproteins, activates Matriptase while Matriptase activates both urokinase-type plasminogen activator and hepatocyte growth factor (HGF). Hepatocyte growth factor activator inhibitor type 1 (HAI-1) is a Kunitz-type serine protease inhibitor identified as a strong inhibitor of Matriptase and HGF. Advanced-stage ovarian tumors that express Matriptase are more likely to do so in the absence of its inhibitor, HAI-1, indicating that an imbalance in the Matriptase:HAI-1 ratio could be important in the development of advanced disease.

## REFERENCES

1. Velasco, G., Cal, S., Quesada, V., Sánchez, L.M. and López-Otín, C. 2002. Matriptase-2, a membrane-bound mosaic serine proteinase predominantly expressed in human liver and showing degrading activity against extracellular matrix proteins. *J. Biol. Chem.* 277: 37637-37646.
2. Hooper, J.D., Campagnolo, L., Goodarzi, G., Truong, T.N., Stuhlmann, H. and Quigley, J.P. 2003. Mouse Matriptase-2: identification, characterization and comparative mRNA expression analysis with mouse Hepsin in adult and embryonic tissues. *Biochem. J.* 373: 689-702.
3. Szabo, R., Netzel-Arnett, S., Hobson, J.P., Antalis, T.M. and Bugge, T.H. 2005. Matriptase-3 is a novel phylogenetically preserved membrane-anchored serine protease with broad serpin reactivity. *Biochem. J.* 390: 231-242.
4. Hartikainen, J.M., Tuhkanen, H., Kataja, V., Eskelinen, M., Uusitupa, M., Kosma, V.M. and Mannermaa, A. 2006. Refinement of the 22q12-q13 breast cancer — associated region: evidence of TMPRSS6 as a candidate gene in an eastern Finnish population. *Clin. Cancer Res.* 12: 1454-1462.
5. Parr, C., Sanders, A.J., Davies, G., Martin, T., Lane, J., Mason, M.D., Mansel, R.E. and Jiang, W.G. 2007. Matriptase-2 inhibits breast tumor growth and invasion and correlates with favorable prognosis for breast cancer patients. *Clin. Cancer Res.* 13: 3568-3576.

## CHROMOSOMAL LOCATION

Genetic locus: Tmprss6 (mouse) mapping to 15 E1.

## PRODUCT

Matriptase-2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Matriptase-2 shRNA Plasmid (m): sc-149299-SH and Matriptase-2 shRNA (m) Lentiviral Particles: sc-149299-V as alternate gene silencing products.

For independent verification of Matriptase-2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149299A, sc-149299B and sc-149299C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Matriptase-2 siRNA (m) is recommended for the inhibition of Matriptase-2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Matriptase-2 gene expression knockdown using RT-PCR Primer: Matriptase-2 (m)-PR: sc-149299-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.