



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

E-Syt1 siRNA (m): sc-149302

BACKGROUND

Synaptotagmins are a large gene family that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. E-Syt1, E-Syt2 and E-Syt3 are Ca^{2+} -regulated intrinsic membrane proteins that belong to the extended synaptotagmins (E-Syts) family. Primary structures and biochemical properties of the E-Syts family are highly conserved from yeast to human and consist of multiple C2 domains, which mediate lipid and calcium binding. Ubiquitously expressed with highest expression in brain, members of the E-Syts family form heteromeric complexes and are thought to play a role in the formation of junctions between endoplasmic reticulum and plasma membrane. E-Syt1 (extended synaptotagmin-like protein 1), also known as MBC2 or FAM62A, is a 1,104 amino acid protein that exists as two alternatively spliced isoforms. The gene encoding E-Syt1 is located on human chromosome 12q13.2.

REFERENCES

- Craxton, M. 2007. Evolutionary genomics of plant genes encoding N-terminal-TM-C2 domain proteins and the similar FAM62 genes and synaptotagmin genes of metazoans. *BMC Genomics* 8: 259.
- Min, S.W., Chang, W.P. and Südhof, T.C. 2007. E-Syts, a family of membrane Ca^{2+} -sensor proteins with multiple C2 domains. *Proc. Natl. Acad. Sci. USA* 104: 3823-3828.
- Craxton, M. 2010. A manual collection of Syt, Esyt, Rph3a, Rph3al, Doc2, and Dblc2 genes from 46 metazoan genomes—an open access resource for neuroscience and evolutionary biology. *BMC Genomics* 11: 37.
- Giordano, F., Saheki, Y., Idevall-Hagren, O., Colombo, S.F., Pirruccello, M., Milosevic, I., Gracheva, E.O., Bagriantsev, S.N., Borgese, N. and De Camilli, P. 2013. PI(4,5)P(2)-dependent and Ca^{2+} -regulated ER-PM interactions mediated by the extended synaptotagmins. *Cell* 153: 1494-1509.
- Herdman, C., Tremblay, M.G., Mishra, P.K. and Moss, T. 2014. Loss of Extended synaptotagmins ESyt2 and ESyt3 does not affect mouse development or viability, but *in vitro* cell migration and survival under stress are affected. *Cell Cycle* 13: 2616-2625.
- Tremblay, M.G., Herdman, C., Guillou, F., Mishra, P.K., Baril, J., Bellenfant, S. and Moss, T. 2015. Extended synaptotagmin interaction with the fibroblast growth factor receptor depends on receptor conformation, not catalytic activity. *J. Biol. Chem.* 290: 16142-16156.
- Föcking, M., Lopez, L.M., English, J.A., Dicker, P., Wolff, A., Brindley, E., Wynne, K., Cagney, G. and Cotter, D.R. 2015. Proteomic and genomic evidence implicates the postsynaptic density in schizophrenia. *Mol. Psychiatry* 20: 424-432.

CHROMOSOMAL LOCATION

Genetic locus: Esyt1 (mouse) mapping to 10 D3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

E-Syt1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see E-Syt1 shRNA Plasmid (m): sc-149302-SH and E-Syt1 shRNA (m) Lentiviral Particles: sc-149302-V as alternate gene silencing products.

For independent verification of E-Syt1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149302A, sc-149302B and sc-149302C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

E-Syt1 siRNA (m) is recommended for the inhibition of E-Syt1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor E-Syt1 gene expression knockdown using RT-PCR Primer: E-Syt1 (m)-PR: sc-149302-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.