



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



ME3 siRNA (m): sc-149344

BACKGROUND

ME3 (malic enzyme 3), also known as NADP-ME (NADP-dependent malic enzyme, mitochondrial), is a 604 amino acid protein of the mitochondrial matrix that catalyzes the conversion of (S)-malate and NADP⁺ to pyruvate, carbon dioxide and NADPH. A member of the malic enzymes family, ME3 utilizes manganese and magnesium as cofactors and is expressed in tissues that divide at a low rate. The gene encoding ME3 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that maps to chromosome 11.

REFERENCES

1. Siebert, G., Ritter, H. and Kömpf, J. 1979. Mitochondrial malic enzyme (E.C.1.1.1.40) in human leukocytes: formal genetics and population genetics. *Hum. Genet.* 51: 319-322.
2. Loeber, G., Maurer-Fogy, I. and Schwendenwein, R. 1994. Purification, cDNA cloning and heterologous expression of the human mitochondrial NADP⁺-dependent malic enzyme. *Biochem. J.* 304: 687-692.
3. Jira, P.E., Waterham, H.R., Wanders, R.J., Smeitink, J.A., Sengers, R.C. and Wevers, R.A. 2003. Smith-Lemli-Opitz syndrome and the DHCR7 gene. *Ann. Hum. Genet.* 67: 269-280.
4. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 604626. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Hsieh, J.Y., Liu, G.Y. and Hung, H.C. 2008. Influential factor contributing to the isoform-specific inhibition by ATP of human mitochondrial NAD(P)⁺-dependent malic enzyme: functional roles of the nucleotide binding site Lys346. *FEBS J.* 275: 5383-5392.
6. Bhuiyan, Z.A., Momenah, T.S., Amin, A.S., Al-Khadra, A.S., Alders, M., Wilde, A.A. and Mannens, M.M. 2008. An intronic mutation leading to incomplete skipping of exon-2 in KCNQ1 rescues hearing in Jervell and Lange-Nielsen syndrome. *Prog. Biophys. Mol. Biol.* 98: 319-327.
7. Hsieh, J.Y., Liu, J.H., Fang, Y.W. and Hung, H.C. 2009. Dual roles of Lys(57) at the dimer interface of human mitochondrial NAD(P)⁺-dependent malic enzyme. *Biochem. J.* 420: 201-209.
8. Hsieh, J.Y., Chen, S.H. and Hung, H.C. 2009. Functional roles of the tetramer organization of malic enzyme. *J. Biol. Chem.* 284: 18096-18105.
9. Coldren, C.D., Lai, Z., Shragg, P., Rossi, E., Glidewell, S.C., Zuffardi, O., Mattina, T., Ivy, D.D., Curfs, L.M., Mattson, S.N., Riley, E.P., Treier, M. and Grossfeld, P.D. 2009. Chromosomal microarray mapping suggests a role for BSX and Neurogranin in neurocognitive and behavioral defects in the 11q terminal deletion disorder (Jacobsen syndrome). *Neurogenetics* 10: 89-95.

CHROMOSOMAL LOCATION

Genetic locus: Me3 (mouse) mapping to 7 E1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

ME3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ME3 shRNA Plasmid (m): sc-149344-SH and ME3 shRNA (m) Lentiviral Particles: sc-149344-V as alternate gene silencing products.

For independent verification of ME3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149344A, sc-149344B and sc-149344C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ME3 siRNA (m) is recommended for the inhibition of ME3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ME3 gene expression knockdown using RT-PCR Primer: ME3 (m)-PR: sc-149344-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.