



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mesogenin 1 siRNA (m): sc-149372

BACKGROUND

Mesogenin 1, also known as MSGN1, pMesogenin1, paraxial mesogenin or MSOG, is a 193 amino acid nuclear protein that plays a role in the formation and segmentation of the paraxial mesoderm. A member of the basic helix-loop-helix (bHLH) family, Mesogenin 1 contains one bHLH domain and is encoded by a gene that maps to human chromosome 2p24.2. Chromosome 2 is the second largest human chromosome and consists of approximately 237 million bases, encodes over 1,400 genes and makes up nearly 8% of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alström syndrome is due to mutations in the ALMS1 gene.

REFERENCES

1. Patel, S.B., Salen, G., Hidaka, H., Kwiterovich, P.O., Stalenhoef, A.F., Miettinen, T.A., Grundy, S.M., Lee, M.H., Rubenstein, J.S., Polymeropoulos, M.H. and Brownstein, M.J. 1998. Mapping a gene involved in regulating dietary cholesterol absorption. The sitosterolemia locus is found at chromosome 2p21. *J. Clin. Invest.* 102: 1041-1044.
2. Yoon, J.K., Moon, R.T. and Wold, B. 2000. The bHLH class protein pMesogenin1 can specify paraxial mesoderm phenotypes. *Dev. Biol.* 222: 376-391.
3. Yoon, J.K. and Wold, B. 2000. The bHLH regulator pMesogenin1 is required for maturation and segmentation of paraxial mesoderm. *Genes Dev.* 14: 3204-3214.
4. Shulenin, S., Schriml, L.M., Remaley, A.T., Fojo, S., Brewer, B., Allikmets, R. and Dean, M. 2001. An ATP-binding cassette gene (ABCG5) from the ABCG (White) gene subfamily maps to human chromosome 2p21 in the region of the sitosterolemia locus. *Cytogenet. Cell Genet.* 92: 204-208.
5. Hearn, T., Renforth, G.L., Spalluto, C., Hanley, N.A., Piper, K., Brickwood, S., White, C., Connolly, V., Taylor, J.F., Russell-Eggitt, I., Bonneau, D., Walker, M. and Wilson, D.I. 2002. Mutation of ALMS1, a large gene with a tandem repeat encoding 47 amino acids, causes Alström syndrome. *Nat. Genet.* 31: 79-83.
6. Kelsell, D.P., Norgett, E.E., Unsworth, H., Teh, M.T., Cullup, T., Mein, C.A., Dopping-Hepenstal, P.J., Dale, B.A., Tadini, G., Fleckman, P., Stephens, K.G., Sybert, V.P., Mallory, S.B., North, B.V., Witt, D.R., Sprecher, E., et al. 2005. Mutations in ABCA12 underlie the severe congenital skin disease harlequin ichthyosis. *Am. J. Hum. Genet.* 76: 794-803.
7. Gomez, C., Ozbudak, E.M., Wunderlich, J., Baumann, D., Lewis, J. and Pourquie, O. 2008. Control of segment number in vertebrate embryos. *Nature* 454: 335-339.

CHROMOSOMAL LOCATION

Genetic locus: Msn1 (mouse) mapping to 12 A1.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Mesogenin 1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Mesogenin 1 shRNA Plasmid (m): sc-149372-SH and Mesogenin 1 shRNA (m) Lentiviral Particles: sc-149372-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Mesogenin 1 siRNA (m) is recommended for the inhibition of Mesogenin 1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Mesogenin 1 gene expression knockdown using RT-PCR Primer: Mesogenin 1 (m)-PR: sc-149372-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.