



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



MIF4GD siRNA (m): sc-149432



The Power to Question

BACKGROUND

MIF4GD (MIF4G domain containing), also known as MIFD, AD023 or SLIP1, is a 222 amino acid protein that contains one MIF4G domain. Localized to both the nucleus and the cytoplasm, MIF4GD plays a role in the replication-dependent translation of histone mRNAs, which differ from most eukaryotic mRNAs in that they end with a stem-loop instead of a poly-A tail. Specifically, MIF4GD interacts with SLBP, eIF4G and DAP-5. Via its interaction with SLBP, MIF4GD is thought to be tethered to the stem loops of histone mRNAs where it may facilitate the circularizing of the mRNAs, thereby enhancing their translation. Depletion of MIF4GD results in reduced histone translation and may lead to cell death, suggesting that MIF4GD plays an important role in cell survival. Two isoforms of MIF4GD exist due to alternative splicing events.

REFERENCES

1. Craig, A.W., Haghigat, A., Yu, A.T. and Sonenberg, N. 1998. Interaction of polyadenylate-binding protein with the eIF4G homologue PAIP enhances translation. *Nature* 392: 520-523.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 612072. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Lehner, B. and Sanderson, C.M. 2004. A protein interaction framework for human mRNA degradation. *Genome Res.* 14: 1315-1323.
4. Rual, J.F., Venkatesan, K., Hao, T., Hirozane-Kishikawa, T., Dricot, A., Li, N., Berriz, G.F., Gibbons, F.D., Dreze, M., Ayivi-Guedehoussou, N., Klitgord, N., Simon, C., Boxem, M., Milstein, S., Rosenberg, J., Goldberg, D.S., et al. 2005. Towards a proteome-scale map of the human protein-protein interaction network. *Nature* 437: 1173-1178.
5. Lim, J., Hao, T., Shaw, C., Patel, A.J., Szabó, G., Rual, J.F., Fisk, C.J., Li, N., Smolyar, A., Hill, D.E., Barabási, A.L., Vidal, M. and Zoghbi, H.Y. 2006. A protein-protein interaction network for human inherited ataxias and disorders of Purkinje cell degeneration. *Cell* 125: 801-814.
6. Hinton, T.M., Coldwell, M.J., Carpenter, G.A., Morley, S.J. and Pain, V.M. 2007. Functional analysis of individual binding activities of the scaffold protein eIF4G. *J. Biol. Chem.* 282: 1695-1708.
7. Cakmakci, N.G., Lerner, R.S., Wagner, E.J., Zheng, L. and Marzluff, W.F. 2008. SLIP1, a factor required for activation of histone mRNA translation by the stem-loop binding protein. *Mol. Cell. Biol.* 28: 1182-1194.

CHROMOSOMAL LOCATION

Genetic locus: Mif4gd (mouse) mapping to 11 E2.

PRODUCT

MIF4GD siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MIF4GD shRNA Plasmid (m): sc-149432-SH and MIF4GD shRNA (m) Lentiviral Particles: sc-149432-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MIF4GD siRNA (m) is recommended for the inhibition of MIF4GD expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MIF4GD gene expression knockdown using RT-PCR Primer: MIF4GD (m)-PR: sc-149432-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.