



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mitsugumin23 siRNA (m): sc-149451

BACKGROUND

With approximately 135 million base pairs and 1,400 genes, chromosome 11 makes up around 4% of human genomic DNA and is considered a gene and disease association dense chromosome. Defects in chromosome 11 have been linked to many diseases, including Wilms' tumors, WAGR syndrome, and Denys-Drash syndrome, Sickle cell anemia and β thalassemia. On chromosome 11q12.2 is a gene encoding Mitsugumin23. Mitsugumin23 (Mg23), also known as transmembrane protein 109 (TMEM109), is a 243 amino acid multi-pass transmembrane protein localized to the membrane of the nucleus, endoplasmic reticulum and sarcoplasmic reticulum. It is comprised of two primary domains: a 33 amino acid signal peptide and a 109 amino acid chain consisting of four transmembrane domains.

REFERENCES

1. Brandt, N.R. and Caswell, A.H. 1999. Localization of mitsugumin 29 to transverse tubules in rabbit skeletal muscle. *Arch. Biochem. Biophys.* 371: 348-350.
2. Nagaraj, R.Y., Nosek, C.M., Brotto, M.A., Nishi, M., Takeshima, H., Nosek, T.M. and Ma, J. 2000. Increased susceptibility to fatigue of slow- and fast-twitch muscles from mice lacking the MG29 gene. *Physiol. Genomics*. 4: 43-49.
3. Brandt, N.R., Franklin, G., Brunschwig, J.P. and Caswell, A.H. 2001. The role of mitsugumin 29 in transverse tubules of rabbit skeletal muscle. *Arch. Biochem. Biophys.* 385: 406-409.
4. Pan, Z., Yang, D., Nagaraj, R.Y., Nosek, T.A., Nishi, M., Takeshima, H., Cheng, H. and Ma, J. 2002. Dysfunction of store-operated calcium channel in muscle cells lacking mg29. *Nat. Cell Biol.* 4: 379-383.
5. Brotto, M.A., Nagaraj, R.Y., Brotto, L.S., Takeshima, H., Ma, J.J. and Nosek, T.M. 2004. Defective maintenance of intracellular Ca^{2+} homeostasis is linked to increased muscle fatigability in the MG29 null mice. *Cell Res.* 14: 373-378.
6. Pan, Z., Hirata, Y., Nagaraj, R.Y., Zhao, J., Nishi, M., Hayek, S.M., Bhat, M.B., Takeshima, H. and Ma, J. 2004. Co-expression of MG29 and ryanodine receptor leads to apoptotic cell death: effect mediated by intracellular Ca^{2+} release. *J. Biol. Chem.* 279: 19387-19390.
7. Weisleder, N., Brotto, M., Komazaki, S., Pan, Z., Zhao, X., Nosek, T., Parness, J., Takeshima, H. and Ma, J. 2006. Muscle aging is associated with compromised Ca^{2+} spark signaling and segregated intracellular Ca^{2+} release. *J. Cell Biol.* 174: 639-645.
8. Zhao, X., Weisleder, N., Thornton, A., Oppong, Y., Campbell, R., Ma, J. and Brotto, M. 2008. Compromised store-operated Ca^{2+} entry in aged skeletal muscle. *Aging Cell* 7: 561-568.
9. Masumiya, H., Asaumi, Y., Nishi, M., Minamisawa, S., Adachi-Akahane, S., Yoshida, M., Kangawa, K., Ito, K., Kagaya, Y., Yanagisawa, T., Yamazaki, T., Ma, J. and Takeshima, H. 2009. Mitsugumin 53-mediated maintenance of K^{+} currents in cardiac myocytes. *Channels* 3: 6-11.

CHROMOSOMAL LOCATION

Genetic locus: Tmem109 (mouse) mapping to 19 A.

PRODUCT

Mitsugumin23 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Mitsugumin23 shRNA Plasmid (m): sc-149451-SH and Mitsugumin23 shRNA (m) Lentiviral Particles: sc-149451-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Mitsugumin23 siRNA (m) is recommended for the inhibition of Mitsugumin23 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Mitsugumin23 gene expression knockdown using RT-PCR Primer: Mitsugumin23 (m)-PR: sc-149451-PR (20 μ l). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.