



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## MYLK3 siRNA (m): sc-149464

### BACKGROUND

The  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinases (CaM kinases) are a structurally related subfamily of serine/threonine kinases that includes CaMKI, CaMKII, CaMKIV and Myosin light chain kinases (MYLKs, also designated MLCs). The MYLK kinases phosphorylate Myosin regulatory light chains to catalyze Myosin interaction with Actin filaments, resulting in contractile activity. MYLK3 (myosin light chain kinase 3) is a 795 amino acid cardiac-specific protein that contains one protein kinase domain and belongs to the protein kinase superfamily. Like other MYLK kinases, MYLK3 is thought to play a role in smooth muscle contraction, specifically using magnesium as a cofactor to catalyze the ATP-dependent phosphorylation of Myosin light chain proteins. Additionally, MYLK3 may regulate sarcomere assembly in heart tissue, possibly mediating proper heart function.

### REFERENCES

1. Lazar, V., et al. 1999. A single human myosin light chain kinase gene (MLCK; MYLK). *Genomics* 57: 256-267.
2. Giorgi, D., et al. 2001. The myosin light chain kinase gene is not duplicated in mouse: partial structure and chromosomal localization of Mylk. *Genomics* 75: 49-56.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 612147. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Seguchi, O., et al. 2007. A cardiac myosin light chain kinase regulates sarcomere assembly in the vertebrate heart. *J. Clin. Invest.* 117: 2812-2824.
5. Chan, J.Y., et al. 2008. Identification of cardiac-specific Myosin light chain kinase. *Circ. Res.* 102: 571-580.
6. Christie, J.D., et al. 2008. Variation in the MYLK gene is associated with development of acute lung injury after major trauma. *Crit. Care Med.* 36: 2794-2800.
7. Al-Sadi, R., et al. 2008. Mechanism of IL-1 $\beta$ -induced increase in intestinal epithelial tight junction permeability. *J. Immunol.* 180: 5653-5661.

### CHROMOSOMAL LOCATION

Genetic locus: Mylk3 (mouse) mapping to 8 C3.

### PRODUCT

MYLK3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MYLK3 shRNA Plasmid (m): sc-149464-SH and MYLK3 shRNA (m) Lentiviral Particles: sc-149464-V as alternate gene silencing products.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}\text{C}$ , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

MYLK3 siRNA (m) is recommended for the inhibition of MYLK3 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MYLK3 gene expression knockdown using RT-PCR Primer: MYLK3 (m)-PR: sc-149464-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^{\circ}\text{C}$  and the extension temperature should be  $68-72^{\circ}\text{C}$ .

### RESEARCH USE

For research use only, not for use in diagnostic procedures.