



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# MOCS3 siRNA (m): sc-149496

## BACKGROUND

Molybdenum is an essential trace element found in most organisms and it functions as a cofactor for several enzymes that catalyze important transformations in carbon, nitrogen and sulfur metabolism. The molybdenum cofactor biosynthetic pathway is evolutionarily conserved between organisms. MOCS3 (molybdenum cofactor synthesis protein 3), also known as UBA4, molybdopterin synthase sulfurylase or MPT synthase sulfurylase, belongs to the hesA/moeB/thiF family and is necessary for the function of all molybdoenzymes. MOCS3 is thought to activate molybdopterin synthase by adenylating its smaller subunit at the C-terminus during molybdopterin biosynthesis in humans. Molybdopterin synthase catalyzes the formation of molybdopterin by incorporating a dithiolene functional group. Molybdenum cofactor deficiency in humans results in the loss of the activity of molybdoenzymes sulfite oxidase, xanthine dehydrogenase and aldehyde oxidase which leads to progressive neurological damage. All forms of molybdenum cofactor deficiency are inherited as autosomal recessive traits.

## REFERENCES

1. Sloan, J., et al. 1999. The two subunits of human molybdopterin synthase: evidence for a bicistronic messenger RNA with overlapping reading frames. *Nucleic Acids Res.* 27: 854-858.
2. Wuebbens, M.M., et al. 2000. Insights into molybdenum cofactor deficiency provided by the crystal structure of the molybdenum cofactor biosynthesis protein MoaC. *Structure* 8: 709-718.
3. Reiss, J. and Johnson, J.L. 2003. Mutations in the molybdenum cofactor biosynthetic genes MOCS1, MOCS2, and GEPH. *Hum. Mutat.* 21: 569-576.
4. Matthies, A., et al. 2004. Evidence for the physiological role of a rhodanese-like protein for the biosynthesis of the molybdenum cofactor in humans. *Proc. Natl. Acad. Sci. USA* 101: 5946-5951.
5. Matthies, A., et al. 2005. Molybdenum cofactor biosynthesis in humans: identification of a persulfide group in the rhodanese-like domain of MOCS3 by mass spectrometry. *Biochemistry* 44: 7912-7920.
6. Leimkühler, S., et al. 2005. Ten novel mutations in the molybdenum cofactor genes MOCS1 and MOCS2 and *in vitro* characterization of a MOCS2 mutation that abolishes the binding ability of molybdopterin synthase. *Hum. Genet.* 117: 565-570.
7. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609277. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Ichida, K., et al. 2006. A Turkish case with molybdenum cofactor deficiency. *Nucleosides Nucleotides Nucleic Acids.* 25: 1087-1091.
9. Krepsinsky, K. and Leimkühler, S. 2007. Site-directed mutagenesis of the active site loop of the rhodanese-like domain of the human molybdopterin synthase sulfurylase MOCS3. Major differences in substrate specificity between eukaryotic and bacterial homologs. *FEBS J.* 274: 2778-2787.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CHROMOSOMAL LOCATION

Genetic locus: Mocs3 (mouse) mapping to 2 H3.

## PRODUCT

MOCS3 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MOCS3 shRNA Plasmid (m): and MOCS3 shRNA (m) Lentiviral Particles: sc-149496-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

MOCS3 siRNA (m) is recommended for the inhibition of MOCS3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MOCS3 gene expression knockdown using RT-PCR Primer: MOCS3 (m)-PR: sc-149496-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.