



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MPHOSPH8 siRNA (m): sc-149530

BACKGROUND

MPHOSPH8 (M-phase phosphoprotein 8), also known as two hybrid-associated protein 3 with RanBPM (Twa), is a 860 amino acid nuclear protein that is phosphorylated in the M (mitotic) phase of the cell cycle. MPHOSPH8 contains four ANK repeats, motifs that are typically found in transcriptional regulators and cell cycle proteins and may be involved in protein-protein binding. Due to evidence suggesting that MPHOSPH8 forms a complex with Muskelein and Ran BP-M, a GTPase implicated in a diverse array of cellular processes, it may play a role in the Ran GTPase cycle. The gene encoding MPHOSPH8 resides on chromosome 13. There are two isoforms of MPHOSPH8 that are produced as a result of alternative splicing events.

REFERENCES

1. Matsumoto-Taniura, N., Pirollet, F., Monroe, R., Gerace, L. and Westendorf, J.M. 1996. Identification of novel M phase phosphoproteins by expression cloning. *Mol. Biol. Cell* 7: 1455-1469.
2. Umeda, M., Nishitani, H. and Nishimoto, T. 2003. A novel nuclear protein, Twa1, and Muskelein comprise a complex with RanBPM. *Gene* 303: 47-54.
3. Kobayashi, N., Yang, J., Ueda, A., Suzuki, T., Tomaru, K., Takeno, M., Okuda, K. and Ishigatsubo, Y. 2007. RanBPM, Muskelein, p48EMLP, p44CTLH, and the armadillo-repeat proteins ARMC8 α and ARMC8 β are components of the CTLH complex. *Gene* 396: 236-247.
4. Voronin, D.A. and Kiseleva, E.V. 2007. Functional role of proteins containing ankyrin repeats. *Tsitologiya* 49: 989-999.
5. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611626. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Valiyaveetil, M., Bentley, A.A., Gursahaney, P., Hussien, R., Chakravarti, R., Kureishy, N., Prag, S. and Adams, J.C. 2008. Novel role of the muskelein-RanBP9 complex as a nucleocytoplasmic mediator of cell morphology regulation. *J. Cell Biol.* 182: 727-739.
7. Dansereau, D.A. and Lasko, P. 2008. RanBPM regulates cell shape, arrangement, and capacity of the female germline stem cell niche in *Drosophila melanogaster*. *J. Cell Biol.* 182: 963-977.
8. Gaudet, R. 2008. A primer on ankyrin repeat function in TRP channels and beyond. *Mol. Biosyst.* 4: 372-379.

CHROMOSOMAL LOCATION

Genetic locus: Mphosph8 (mouse) mapping to 14 C3.

PRODUCT

MPHOSPH8 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MPHOSPH8 shRNA Plasmid (m): sc-149530-SH and MPHOSPH8 shRNA (m) Lentiviral Particles: sc-149530-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MPHOSPH8 siRNA (m) is recommended for the inhibition of MPHOSPH8 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

MPHOSPH8 (C-8): sc-398598 is recommended as a control antibody for monitoring of MPHOSPH8 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MPHOSPH8 gene expression knockdown using RT-PCR Primer: MPHOSPH8 (m)-PR: sc-149530-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.