



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MPPE1 siRNA (m): sc-149539

BACKGROUND

Metallophosphoesterases (MPPEs) are dynamic enzymes that catalyze a variety of cellular reactions and contain a conserved β - α - β - α fold. The MPPE superfamily is divided into two subfamilies: phosphomonoesterases and phosphodiesterases. Each MPPE has a dimetal center located approximately at the C-terminal end of the parallel β -strands of the fold. MPPE1 (metallophosphoesterase 1) is a 396 amino acid multipass membrane enzyme that requires two divalent metals as cofactors. MPPE1 contains a N-terminal signal peptide, a typical metallophosphoesterase domain and a C-terminal transmembrane domain. Expression of MPPE1 seems to be limited to brain. There are five isoforms of MPPE1 that are produced as a result of alternative splicing events.

REFERENCES

- Chern, T.H., Chiang, F.T., Hsu, K.L., Lo, H.M., Tseng, C.D. and Tseng, Y.Z. 2000. Cloning a novel metallophosphoesterase gene from a kidney cDNA library of hypertensive rat. *J. Formos. Med. Assoc.* 99: 49-53.
- Vuoristo, J.T. and Ala-Kokko, L. 2001. cDNA cloning, genomic organization and expression of the novel human metallophosphoesterase gene MPPE1 on chromosome 18p11.2. *Cytogenet. Cell Genet.* 95: 60-63.
- Wan, D., Gong, Y., Qin, W., Zhang, P., Li, J., Wei, L., Zhou, X., Li, H., Qiu, X., Zhong, F., He, L., Yu, J., Yao, G., Jiang, H., Qian, L., Yu, Y., Shu, H., Chen, X., Xu, H., Guo, M., Pan, Z., Chen, Y., Ge, C., Yang, S. and Gu, J. 2004. Large-scale cDNA transfection screening for genes related to cancer development and progression. *Proc. Natl. Acad. Sci. USA* 101: 15724-15729.
- Keppetipola, N. and Shuman, S. 2006. Distinct enzymic functional groups are required for the phosphomonoesterase and phosphodiesterase activities of *Clostridium thermocellum* polynucleotide kinase/phosphatase. *J. Biol. Chem.* 281: 19251-19259.
- Keppetipola, N. and Shuman, S. 2007. Characterization of the 2',3' cyclic phosphodiesterase activities of *Clostridium thermocellum* polynucleotide kinase-phosphatase and bacteriophage lambda phosphatase. *Nucleic Acids Res.* 35: 7721-7732.
- Canales, J., Fernández, A., Ribeiro, J.M., Cabezas, A., Rodrigues, J.R., Cameselle, J.C. and Costas, M.J. 2008. Mn²⁺-dependent ADP-ribose/CDP-alcohol pyrophosphatase: a novel metallophosphoesterase family preferentially expressed in rodent immune cells. *Biochem. J.* 413: 103-113.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 611900. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Tyagi, R., Shenoy, A.R. and Visweswariah, S.S. 2009. Characterization of an evolutionarily conserved metallophosphoesterase that is expressed in the fetal brain and associated with the WAGR syndrome. *J. Biol. Chem.* 284: 5217-5228.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Mppe1* (mouse) mapping to 18 E1.

PRODUCT

MPPE1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MPPE1 shRNA Plasmid (m): sc-149539-SH and MPPE1 shRNA (m) Lentiviral Particles: sc-149539-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MPPE1 siRNA (m) is recommended for the inhibition of MPPE1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MPPE1 gene expression knockdown using RT-PCR Primer: MPPE1 (m)-PR: sc-149539-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.