



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CLEC-18 siRNA (m): sc-149549

BACKGROUND

The C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily consists of a variety of proteins that share a common protein fold and have diverse functions, including cell-cell signaling, cell adhesion, glycoprotein turnover and immune responses. CLEC-18 (C-type lectin domain family 18) exists as three nearly identical forms, designated CLEC-18A, CLEC-18B and CLEC-18C, or mannose receptor-like protein 1, 2 and 3, respectively. Both CLEC-18A and CLEC-18C contain 446 amino acids, whereas CLEC-18B contains 455 amino acids; all 3 forms are secreted proteins that contain one C-type lectin domain, a SCP domain and a single EGF-like domain. All three CLEC-18 proteins exist as two alternatively spliced isoforms that are encoded by genes that map to human chromosome 16q22.

REFERENCES

1. Drickamer, K. 1999. C-type lectin-like domains. *Curr. Opin. Struct. Biol.* 9: 585-590.
2. Arce, I., et al. 2001. Molecular and genomic characterization of human DLEC, a novel member of the C-type lectin receptor gene family preferentially expressed on monocyte-derived dendritic cells. *Eur. J. Immunol.* 31: 2733-2740.
3. East, L., et al. 2002. The mannose receptor family. *Biochim. Biophys. Acta* 1572: 364-386.
4. Ebner, S., et al. 2003. Evolutionary analysis reveals collective properties and specificity in the C-type lectin and lectin-like domain superfamily. *Proteins* 53: 44-55.
5. McMahon, S.A., et al. 2005. The C-type lectin fold as an evolutionary solution for massive sequence variation. *Nat. Struct. Mol. Biol.* 12: 886-892.

CHROMOSOMAL LOCATION

Genetic locus: Clec18a (mouse) mapping to 8 E1.

PRODUCT

CLEC-18 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CLEC-18 shRNA Plasmid (m): sc-149549-SH and CLEC-18 shRNA (m) Lentiviral Particles: sc-149549-V as alternate gene silencing products.

For independent verification of CLEC-18 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149549A, sc-149549B and sc-149549C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CLEC-18 siRNA (m) is recommended for the inhibition of CLEC-18 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CLEC-18 gene expression knockdown using RT-PCR Primer: CLEC-18 (m)-PR: sc-149549-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.