



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



MRS2L siRNA (m): sc-149639

BACKGROUND

MRS2L (magnesium transporter MRS2 homolog, mitochondrial), also known as MRS2 or HPT, is a 443 amino acid protein that is a part of the inner membrane of mitochondrion and is distantly related to CorA Mg²⁺ transport proteins. Belonging to the CorA metal ion transporter (MIT) (TC 1.A.35) family, MRS2L is a magnesium transporter that may mediate the influx of magnesium into the mitochondrial matrix. MRS2L exists as three alternatively spliced isoforms and is considered a multi-pass membrane protein. MRS2L, along with OXA1L, YME1L1 and MIPEP, may be involved in the pathology of combined enzymatic deficiencies of the oxidative phosphorylation (OXPHOS) system. MRS2L deregulation can promote cell death because deficiency of external magnesium induces apoptosis in rat hepatocytes. In brain MRS2L is most highly expressed in the superior temporal cortex, hypothalamus and amygdala. The MRS2L gene maps to chromosome 6p22.2.

REFERENCES

1. London, E.R., Meng, H. and Gruen, J.R. 2003. A transcription map of the 6p22.3 reading disability locus identifying candidate genes. *BMC Genomics* 4: 25.
2. Cope, N., Harold, D., Hill, G., Moskvina, V., Stevenson, J., Holmans, P., Owen, M.J., O'Donovan, M.C. and Williams, J. 2005. Strong evidence that KIAA0319 on chromosome 6p is a susceptibility gene for developmental dyslexia. *Am. J. Hum. Genet.* 76: 581-591.
3. Younossi, Z.M., Baranova, A., Ziegler, K., Del Giacco, L., Schlauch, K., Born, T.L., Elariny, H., Gorreta, F., VanMeter, A., Younoszai, A., Ong, J.P., Goodman, Z. and Chandhoke, V. 2005. A genomic and proteomic study of the spectrum of nonalcoholic fatty liver disease. *Hepatology* 42: 665-674.
4. Coenen, M.J., Smeitink, J.A., Smeets, R., Trijbels, F.J. and van den Heuvel, L.P. 2005. Mutation detection in four candidate genes (OXA1L, MRS2L, YME1L and MIPEP) for combined deficiencies in the oxidative phosphorylation system. *J. Inherit. Metab. Dis.* 28: 1091-1097.
5. Meng, H., Smith, S.D., Hager, K., Held, M., Liu, J., Olson, R.K., Pennington, B.F., DeFries, J.C., Gelernter, J., O'Reilly-Pol, T., Somlo, S., Skudlarski, P., Shaywitz, S.E., Shaywitz, B.A., Marchione, K., Wang, Y., et al. 2005. DCDC2 is associated with reading disability and modulates neuronal development in the brain. *Proc. Natl. Acad. Sci. USA* 102: 17053-17058.
6. Piskacek, M., Zotova, L., Zsurka, G. and Schweny, R.J. 2009. Conditional knockdown of hMRS2 results in loss of mitochondrial Mg²⁺ uptake and cell death. *J. Cell. Mol. Med.* 13: 693-700.
7. Martin, R.E., Ginsburg, H. and Kirk, K. 2009. Membrane transport proteins of the malaria parasite. *Mol. Microbiol.* 74: 519-528.
8. SWISS-PROT/TrEMBL (Q9HD23). World Wide Web URL: <http://www.uniprot.org/uniprot/Q9HD23>

CHROMOSOMAL LOCATION

Genetic locus: Mrs2 (mouse) mapping to 13 A3.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

MRS2L siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MRS2L shRNA Plasmid (m): sc-149639-SH and MRS2L shRNA (m) Lentiviral Particles: sc-149639-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MRS2L siRNA (m) is recommended for the inhibition of MRS2L expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MRS2L gene expression knockdown using RT-PCR Primer: MRS2L (m)-PR: sc-149639-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.