



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MYL6 siRNA (m): sc-149750

BACKGROUND

The EF-hand domain is a twelve amino acid loop motif that is commonly found in proteins that participate in calcium-binding events within the cell. EF-hand domains generally exist in a pair that, together, form a stable four-helix bundle that enables the binding of calcium ions. MYL6 (myosin, light chain 6, alkali, smooth muscle and non-muscle), also known as ESMC, LC17A, LC17B or MLC1SM, is a 151 amino acid protein that contains three EF-hand domains and exists as two alternatively spliced isoforms, designated smooth muscle (MLC3SM) and non-muscle (MLC3NM). Existing as an alkali light chain component of the hexameric myosin complex, MYL6 participates in generating the force for cellular movements, thereby playing an important role in overall cellular function. The gene encoding MYL6 maps to human chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5% of the human genome.

REFERENCES

1. Lenz, S., Lohse, P., Seidel, U. and Arnold, H.H. 1989. The alkali light chains of human smooth and nonmuscle myosins are encoded by a single gene. Tissue-specific expression by alternative splicing pathways. *J. Biol. Chem.* 264: 9009-9015.
2. Hailstones, D.L. and Gunning, P.W. 1990. Characterization of human myosin light chains 1sa and 3nm: implications for isoform evolution and function. *Mol. Cell. Biol.* 10: 1095-1104.
3. Bora, P.S., Bora, N.S., Wu, X., Kaplan, H.J. and Lange, L.G. 1994. Molecular cloning, sequencing, and characterization of smooth muscle myosin alkali light chain from human eye cDNA: homology with myocardial fatty acid ethyl ester synthase-III cDNA. *Genomics* 19: 186-188.
4. Komiyama, M., Soldati, T., von Arx, P. and Perriard, J.C. 1996. The intra-compartmental sorting of myosin alkali light chain isoproteins reflects the sequence of developmental expression as determined by double epitope-tagging competition. *J. Cell Sci.* 109: 2089-2099.
5. Watanabe, M., Kohri, M., Takaishi, M., Horie, R. and Higashihara, M. 2001. Molecular cloning and sequencing of myosin light chains in human megakaryoblastic leukemia cells. *J. Smooth Muscle Res.* 37: 25-38.
6. Fu, Z.Y., Xie, B.T., Ma, Y.T. and Gong, Z.X. 2006. Preparation of monoclonal antibodies against human ventricular myosin light chain 1 (HVMLC1) for functional studies. *Acta Biochim. Biophys. Sin.* 38: 625-632.
7. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609931. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Myl6 (mouse) mapping to 10 D3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

MYL6 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MYL6 shRNA Plasmid (m): sc-149750-SH and MYL6 shRNA (m) Lentiviral Particles: sc-149750-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MYL6 siRNA (m) is recommended for the inhibition of MYL6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MYL6 gene expression knockdown using RT-PCR Primer: MYL6 (m)-PR: sc-149750-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.