



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NCKX6 siRNA (m): sc-149857

BACKGROUND

NCKX6 (Na⁺/K⁺/Ca²⁺-exchange protein 6), also designated SLC24A6 (solute carrier family 24 member 6), belongs to a family of potassium-dependent sodium/calcium exchangers, all of which contain 2 large hydrophilic loops and 2 sets of multiple transmembrane-spanning segments. Potassium-dependent sodium/calcium exchangers maintain cellular calcium homeostasis via the countertransport of four sodium ions for one calcium ion and a potassium ion. Belonging to the SLC24A subfamily and consisting of 584 amino acids, NCKX6 is unlike other potassium-dependent sodium/calcium exchangers in that it functions independently of potassium and transports calcium in exchange for either lithium or sodium. NCKX6 is a multi-pass membrane protein that is strongly inhibited by zinc, exists as two alternatively spliced isoforms and is encoded by a gene mapping to human chromosome 12q24.13.

REFERENCES

1. Lytton, J., et al. 2002. K⁺-dependent Na⁺/Ca²⁺ exchangers in the brain. *Ann. N.Y. Acad. Sci.* 976: 382-393.
2. Cai, X., et al. 2004. Molecular cloning of a sixth member of the K⁺-dependent Na⁺/Ca²⁺ exchanger gene family, NCKX6. *J. Biol. Chem.* 279: 5867-5876.
3. Palty, R., et al. 2004. Lithium-calcium exchange is mediated by a distinct potassium-independent sodium-calcium exchanger. *J. Biol. Chem.* 279: 25234-25240.
4. Schnetkamp, P.P. 2004. The SLC24 Na⁺/Ca²⁺-K⁺ exchanger family: vision and beyond. *Pflugers Arch.* 447: 683-688.
5. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 609841. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Altimimi, H.F., et al. 2007. Examining Ca²⁺ extrusion of Na⁺/Ca²⁺-K⁺ exchangers. *Ann. N.Y. Acad. Sci.* 1099: 29-33.
7. Lytton, J. 2007. Na⁺/Ca²⁺ exchangers: three mammalian gene families control Ca²⁺ transport. *Biochem. J.* 406: 365-382.

CHROMOSOMAL LOCATION

Genetic locus: Slc24a6 (mouse) mapping to 5 F.

PRODUCT

NCKX6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NCKX6 shRNA Plasmid (m): sc-149857-SH and NCKX6 shRNA (m) Lentiviral Particles: sc-149857-V as alternate gene silencing products.

For independent verification of NCKX6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149857A, sc-149857B and sc-149857C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NCKX6 siRNA (m) is recommended for the inhibition of NCKX6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NCKX6 gene expression knockdown using RT-PCR Primer: NCKX6 (m)-PR: sc-149857-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.