



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# NDUFA11 siRNA (m): sc-149868

## BACKGROUND

NDUFA11 (NADH-ubiquinone oxidoreductase a subunit 11), also designated complex I-B14.7, is one of 45 subunits comprising complex I of the oxidative phosphorylation electron transport chain. The multi-subunit NADH:ubiquinone oxidoreductase (complex I) is the first enzyme complex in the electron transport chain of the mitochondria. Complex I deficiency is the most common respiratory chain defect, resulting in various combinations of cardiac, hepatic, and renal disorders. Through use of chaotropic agents, complex I can be separated into three different fractions: a flavoprotein fraction, a hydrophobic protein (HP) fraction and an iron-sulfur protein (IP) fraction. Mutation in the NDUFA11 gene leads to complex I deficiency presenting with infantile lactic acidemia or encephalomyopathy.

## REFERENCES

1. Chow, W., et al. 1991. Determination of the cDNA sequence for the human mitochondrial 75-kDa Fes protein of NADH-coenzyme Q reductase. *Eur. J. Biochem.* 201: 547-550.
2. Duncan, A.M., et al. 1992. Localization of the human 75-kDa Fes protein of NADH-coenzyme Q reductase gene (NDUFS1) to 2q33-q34. *Cytogenet. Cell Genet.* 60: 212-213.
3. Stojanovski, D., et al. 2004. Levels of human Fis1 at the mitochondrial outer membrane regulate mitochondrial morphology. *J. Cell Sci.* 117: 1201-1210.
4. Karahan, O.I., et al. 2005. Ultrasound evaluation of peritoneal catheter tunnel in catheter related infections in CAPD. *Int. Urol. Nephrol.* 37: 363-366.
5. Martin, M.A., et al. 2005. Leigh syndrome associated with mitochondrial complex I deficiency due to a novel mutation in the NDUFS1 gene. *Arch. Neurol.* 62: 659-661.
6. Smeitink, J.A., et al. 2005. Cell biological consequences of mitochondrial NADH: ubiquinone oxidoreductase deficiency. *Curr. Neurovasc. Res.* 1: 29-40.

## CHROMOSOMAL LOCATION

Genetic locus: Ndufa11 (mouse) mapping to 17 D.

## PRODUCT

NDUFA11 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NDUFA11 shRNA Plasmid (m): sc-149868-SH and NDUFA11 shRNA (m) Lentiviral Particles: sc-149868-V as alternate gene silencing products.

For independent verification of NDUFA11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149868A and sc-149868B.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

NDUFA11 siRNA (m) is recommended for the inhibition of NDUFA11 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NDUFA11 gene expression knockdown using RT-PCR Primer: NDUFA11 (m)-PR: sc-149868-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.