



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NDUFB5 siRNA (m): sc-149882

BACKGROUND

Complex 1 (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex 1 is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFB5 (NADH dehydrogenase (ubiquinone) 1 β subcomplex, 5), also known as SGDHI or CI-SGDH, is a single-pass membrane protein that localizes to the mitochondrial inner membrane and functions as an accessory subunit of Complex I.

REFERENCES

1. Ton, C., Hwang, D.M., Dempsey, A.A. and Liew, C.C. 1997. Identification and primary structure of five human NADH-ubiquinone oxidoreductase subunits. *Biochem. Biophys. Res. Commun.* 241: 589-594.
2. Hirst, J., Carroll, J., Fearnley, I.M., Shannon, R.J. and Walker, J.E. 2003. The nuclear encoded subunits of Complex I from bovine heart mitochondria. *Biochim. Biophys. Acta* 1604: 135-150.
3. Sparks, L.M., Xie, H., Koza, R.A., Mynatt, R., Hulver, M.W., Bray, G.A. and Smith, S.R. 2005. A high-fat diet coordinately downregulates genes required for mitochondrial oxidative phosphorylation in skeletal muscle. *Diabetes* 54: 1926-1933.
4. Janssen, R.J., Nijtmans, L.G., van den Heuvel, L.P. and Smeitink, J.A. 2006. Mitochondrial Complex I: structure, function and pathology. *J. Inher. Metab. Dis.* 29: 499-515.
5. Parikh, H., Nilsson, E., Ling, C., Poulsen, P., Almgren, P., Nittby, H., Eriksson, K.F., Vaag, A. and Groop, L.C. 2008. Molecular correlates for maximal oxygen uptake and type 1 fibers. *Am. J. Physiol. Endocrinol. Metab.* 294: E1152-E1159.
6. De Rasmio, D., Panelli, D., Sardanelli, A.M. and Papa, S. 2008. cAMP-dependent protein kinase regulates the mitochondrial import of the nuclear encoded NDUFS4 subunit of Complex I. *Cell. Signal.* 20: 989-997.
7. Lund, K.C. and Wallace, K.B. 2008. Adenosine 3',5'-cyclic monophosphate (cAMP)-dependent phosphoregulation of mitochondrial Complex I is inhibited by nucleoside reverse transcriptase inhibitors. *Toxicol. Appl. Pharmacol.* 226: 94-106.

CHROMOSOMAL LOCATION

Genetic locus: *Ndufb5* (mouse) mapping to 3 A3.

PRODUCT

NDUFB5 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NDUFB5 shRNA Plasmid (m): sc-149882-SH and NDUFB5 shRNA (m) Lentiviral Particles: sc-149882-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NDUFB5 siRNA (m) is recommended for the inhibition of NDUFB5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NDUFB5 (F-2): sc-514245 is recommended as a control antibody for monitoring of NDUFB5 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NDUFB5 gene expression knockdown using RT-PCR Primer: NDUFB5 (m)-PR: sc-149882-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.