



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## NELL1 siRNA (m): sc-149910

### BACKGROUND

NELL1 (NEL-like 1), also known as NRP1 or IDH3GL (NAD<sup>+</sup>-dependent isocitrate dehydrogenase 3  $\gamma$ -like), is an 810 amino acid secreted protein that contains one TSP N-terminal domain, five VWFC domains and six EGF-like domains. Expressed in craniofacial anomalies, NELL1 exists as a homotrimer that interacts with PKC  $\beta$  and is involved in the regulation of cell growth and differentiation. Human NELL1 shares 50% amino acid identity with its chicken counterpart, suggesting that NELL1 may have a conserved role between species. The gene encoding NELL1 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are associated with defects in genes that maps to chromosome 11.

### REFERENCES

1. Watanabe, T.K., et al. 1996. Cloning and characterization of two novel human cDNAs (NELL1 and NELL2) encoding proteins with six EGF-like repeats. *Genomics* 38: 273-276.
2. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 602319: World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Luce, M.J., et al. 1999. The neuronal EGF-related genes NELL1 and NELL2 are expressed in hemopoietic cells and developmentally regulated in the B lineage. *Gene* 231: 121-126.
4. Ting, K., et al. 1999. Human NELL-1 expressed in unilateral coronal synostosis. *J. Bone Miner. Res.* 14: 80-89.
5. Maeda, K., et al. 2001. Brain specific human genes, NELL1 and NELL2, are predominantly expressed in neuroblastoma and other embryonal neuroepithelial tumors. *Neurol. Med. Chir.* 41: 582-588.
6. Okamoto, K., et al. 2003. Identification of NAD<sup>+</sup>-dependent isocitrate dehydrogenase 3  $\gamma$ -like (IDH3GL) gene and its genetic polymorphisms. *Gene* 323: 141-148.

### CHROMOSOMAL LOCATION

Genetic locus: Nell1 (mouse) mapping to 7 B5.

### PRODUCT

NELL1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NELL1 shRNA Plasmid (m): sc-149910-SH and NELL1 shRNA (m) Lentiviral Particles: sc-149910-V as alternate gene silencing products.

For independent verification of NELL1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149910A, sc-149910B and sc-149910C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

NELL1 siRNA (m) is recommended for the inhibition of NELL1 expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NELL1 gene expression knockdown using RT-PCR Primer: NELL1 (m)-PR: sc-149910-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.