



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



NKAP siRNA (m): sc-149988

BACKGROUND

NF κ B, a pleiotropic transcription factor, is present in almost all cell types and is involved in many biological processes including inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF κ B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NF κ B1/p105, NF κ B1/p50, REL and NF κ B2/p52. This complex is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NKAP (NF κ B-activating protein) is a 415 amino acid nuclear protein that regulates TNF and IL1-induced NF κ B activation. As component of a DNA-binding complex, NKAP also functions as a transcriptional repressor that acts on NOTCH target genes. Loss of NKAP blocks the development of $\alpha\beta$ T-cells, suggesting that it is required for their maturation through repression of NOTCH genes.

REFERENCES

1. Bowie, A. and O'Neill, L.A. 1997. Studies into the mechanism of NF κ B activation by IL1, TNF and H₂O₂ in primary and transformed endothelial cells. *Biochem. Soc. Trans.* 25: 125S.
2. Chen, D., Li, X., Zhai, Z. and Shu, H.B. 2002. A novel zinc finger protein interacts with receptor-interacting protein (RIP) and inhibits tumor necrosis factor (TNF)- and IL1-induced NF κ B activation. *J. Biol. Chem.* 277: 15985-15991.
3. Chen, D., Li, Z., Yang, Q., Zhang, J., Zhai, Z. and Shu, H.B. 2003. Identification of a nuclear protein that promotes NF κ B activation. *Biochem. Biophys. Res. Commun.* 310: 720-724.
4. Dai, S.M., Nishioka, K. and Yudoh, K. 2004. Interleukin (IL) 18 stimulates osteoclast formation through synovial T cells in rheumatoid arthritis: comparison with IL1 β and tumour necrosis factor α . *Ann. Rheum. Dis.* 63: 1379-1386.
5. Borggreffe, T. and Oswald, F. 2009. The Notch signaling pathway: transcriptional regulation at Notch target genes. *Cell. Mol. Life Sci.* 66: 1631-1646.
6. Pajeroski, A.G., Nguyen, C., Aghajanian, H., Shapiro, M.J. and Shapiro, V.S. 2009. NKAP is a transcriptional repressor of Notch signaling and is required for T cell development. *Immunity* 30: 696-707.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 300766. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Nkap (mouse) mapping to X A3.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

NKAP siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NKAP shRNA Plasmid (m): sc-149988-SH and NKAP shRNA (m) Lentiviral Particles: sc-149988-V as alternate gene silencing products.

For independent verification of NKAP (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-149988A, sc-149988B and sc-149988C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NKAP siRNA (m) is recommended for the inhibition of NKAP expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NKAP gene expression knockdown using RT-PCR Primer: NKAP (m)-PR: sc-149988-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.