



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



NMB siRNA (m): sc-150007

BACKGROUND

NMB (neuromedin-B) is a 121 amino acid secreted protein that belongs to the bombesin/neuromedin-B/ranatensin family and exists as two alternatively spliced isoforms. Acting in a manner similar to that of bombesin, NMB stimulates smooth muscle contraction. The gene that encodes NMB consists of about 3,443 bases and maps to human chromosome 15q25. Housing approximately 106 million base pairs and encoding more than 700 genes, chromosome 15 makes up about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

REFERENCES

1. Krane, I.M., Naylor, S.L., Helin-Davis, D., Chin, W.W. and Spindel, E.R. 1988. Molecular cloning of cDNAs encoding the human bombesin-like peptide neuromedin B. Chromosomal localization and comparison to cDNAs encoding its amphibian homolog ranatensin. *J. Biol. Chem.* 263: 13317-13323.
2. Online Mendelian Inheritance in Man, OMIM[™]. 1988. Johns Hopkins University, Baltimore, MD. MIM Number: 162340. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Fathi, Z., Benya, R.V., Shapira, H., Jensen, R.T. and Battey, J.F. 1993. The fifth transmembrane segment of the neuromedin B receptor is critical for high affinity neuromedin B binding. *J. Biol. Chem.* 268: 14622-14626.
4. Lee, S. and Kim, Y. 1999. Solution structure of neuromedin B by ¹H nuclear magnetic resonance spectroscopy. *FEBS Lett.* 460: 263-269.
5. Bouchard, L., Drapeau, V., Provencher, V., Lemieux, S., Chagnon, Y., Rice, T., Rao, D.C., Vohl, M.C., Tremblay, A., Bouchard, C. and Perusse, L. 2004. Neuromedin β : a strong candidate gene linking eating behaviors and susceptibility to obesity. *Am. J. Clin. Nutr.* 80: 1478-1486.
6. Kopan, S., Sivasubramanian, U. and Warburton, M.J. 2004. The lysosomal degradation of neuromedin B is dependent on tripeptidyl peptidase-I: evidence for the impairment of neuropeptide degradation in late-infantile neuronal ceroid lipofuscinosis. *Biochem. Biophys. Res. Commun.* 319: 58-65.
7. Hoggard, N., Bashir, S., Cruickshank, M., Miller, J.D. and Speakman, J.R. 2007. Expression of neuromedin B in adipose tissue and its regulation by changes in energy balance. *J. Mol. Endocrinol.* 39: 199-210.
8. Spálová, J., Zamrazilová, H., Vcelák, J., Vanková, M., Lukášová, P., Hill, M., Hlavatá, K., Srámková, P., Fried, M., Aldhoon, B., Kunesová, M., Bendlová, B. and Hainer, V. 2008. Neuromedin β : P73T polymorphism in overweight and obese subjects. *Physiol. Res.* 57: S39-S48.
9. Pigeyre, M., Bokor, S., Romon, M., Gottrand, F., Gilbert, C.C., Valtueña, J., Gómez-Martínez, S., Moreno, L.A., Amouyel, P., Dallongeville, J. and Meirhaeghe, A. 2010. Influence of maternal educational level on the association between the rs3809508 neuromedin B gene polymorphism and the risk of obesity in the HELENA study. *Int. J. Obes.* 34: 478-486.

CHROMOSOMAL LOCATION

Genetic locus: Nmb (mouse) mapping to 7 D3.

PRODUCT

NMB siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NMB shRNA Plasmid (m): sc-150007-SH and NMB shRNA (m) Lentiviral Particles: sc-150007-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NMB siRNA (m) is recommended for the inhibition of NMB expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NMB gene expression knockdown using RT-PCR Primer: NMB (m)-PR: sc-150007-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.