



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NPEPL1 siRNA (m): sc-150049

BACKGROUND

Aminopeptidases are widely distributed in eukaryotes and prokaryotes. These enzymes catalyze the removal of amino acids from the N-termini of proteins or peptide substrates. Aminopeptidases are involved in regulating hormone levels and are essential for digestive and intracellular protein metabolism. NPEPL1, also referred to as aminopeptidase-like 1, is a 523 amino acid protein that belongs to the peptidase M17 family and is ubiquitously expressed. NPEPL1 may be involved in the processing, catabolism and degradation of intracellular proteins by catalyzing the removal of unsubstituted N-terminal amino acids from various peptides. NPEPL1 contains several zinc binding sites and is expressed as three isoforms due to alternative splicing events.

REFERENCES

1. Taylor, A. 1993. Amino peptidases: structure and function. *FASEB J.* 7: 290-298.
2. Taylor, A. 1993. Amino peptidases: towards a mechanism of action. *Trends Biochem. Sci.* 18: 167-171.
3. Meinel, T., Serero, A. and Giglione, C. 2006. Impact of the N-terminal amino acid on targeted protein degradation. *Biol. Chem.* 387: 839-851.
4. Herrera-Camacho, I., Rosas-Murrieta, N.H., Rojo-Domínguez, A., Millán, L., Reyes-Leyva, J., Santos-López, G. and Suárez-Rendueles, P. 2007. Biochemical characterization and structural prediction of a novel cytosolic leucyl aminopeptidase of the M17 family from *Schizosaccharomyces pombe*. *FEBS J.* 274: 6228-6240.
5. Chen, S.L., Marino, T., Fang, W.H., Russo, N. and Himo, F. 2008. Peptide hydrolysis by the binuclear zinc enzyme aminopeptidase from *Aeromonas proteolytica*: a density functional theory study. *J. Phys. Chem. B* 112: 2494-2500.
6. SWISS-PROT/TrEMBL (Q8NDH3). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: Npepl1 (mouse) mapping to 2 H4.

PRODUCT

NPEPL1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NPEPL1 shRNA Plasmid (m): sc-150049-SH and NPEPL1 shRNA (m) Lentiviral Particles: sc-150049-V as alternate gene silencing products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NPEPL1 siRNA (m) is recommended for the inhibition of NPEPL1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NPEPL1 (E-10): sc-515463 is recommended as a control antibody for monitoring of NPEPL1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NPEPL1 gene expression knockdown using RT-PCR Primer: NPEPL1 (m)-PR: sc-150049-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.