



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NPT2c siRNA (m): sc-150057

BACKGROUND

NPT2c (Na⁺-dependent phosphate cotransporter 2C), also known as SLC34A3 (solute carrier family 34 member 3), is a 599 amino acid multi-pass membrane protein that maintains inorganic phosphate concentration at the kidney by assisting in the active transport of phosphate through the renal brush border membrane. A member of the SLC34A transporter family, NPT2c is encoded by a gene that maps to human chromosome 9q34.3. Defects in the NPT2c gene are the cause of hereditary hypophosphatemic rickets with hypercalciuria (HHRH), an autosomal recessive disease characterized by rickets, hypophosphatemia and decreased renal phosphate reabsorption.

REFERENCES

1. Tieder, M., Modai, D., Samuel, R., Arie, R., Halabe, A., Bab, I., Gabizon, D. and Liberman, U.A. 1985. Hereditary hypophosphatemic rickets with hypercalciuria. *N. Engl. J. Med.* 312: 611-617.
2. Segawa, H., Kaneko, I., Takahashi, A., Kuwahata, M., Ito, M., Ohkido, I., Tatsumi, S. and Miyamoto, K. 2002. Growth-related renal type II Na/P_i cotransporter. *J. Biol. Chem.* 277: 19665-19672.
3. Bergwitz, C., Roslin, N.M., Tieder, M., Loredó-Osti, J.C., Bastepe, M., Abu-Zahra, H., Frappier, D., Burkett, K., Carpenter, T.O., Anderson, D., Garabedian, M., Sermet, I., Fujiwara, T.M., Morgan, K., et al. 2006. SLC34A3 mutations in patients with hereditary hypophosphatemic rickets with hypercalciuria predict a key role for the sodium-phosphate cotransporter NaP_i-IIc in maintaining phosphate homeostasis. *Am. J. Hum. Genet.* 78: 179-192.
4. Lorenz-Depiereux, B., Benet-Pages, A., Eckstein, G., Tenenbaum-Rakover, Y., Wagenstaller, J., Tiosano, D., Gershoni-Baruch, R., Albers, N., Lichtner, P., Schnabel, D., Hochberg, Z. and Strom, T.M. 2006. Hereditary hypophosphatemic rickets with hypercalciuria is caused by mutations in the sodium-phosphate cotransporter gene SLC34A3. *Am. J. Hum. Genet.* 78: 193-201.
5. Ichikawa, S., Sorenson, A.H., Imel, E.A., Friedman, N.E., Gertner, J.M. and Econs, M.J. 2006. Intronic deletions in the SLC34A3 gene cause hereditary hypophosphatemic rickets with hypercalciuria. *J. Clin. Endocrinol. Metab.* 91: 4022-4027.
6. Forster, I.C., Hernando, N., Biber, J. and Murer, H. 2006. Proximal tubular handling of phosphate: a molecular perspective. *Kidney Int.* 70: 1548-1559.
7. Yamamoto, T., Michigami, T., Aranami, F., Segawa, H., Yoh, K., Nakajima, S., Miyamoto, K. and Ozono, K. 2007. Hereditary hypophosphatemic rickets with hypercalciuria: a study for the phosphate transporter gene type IIc and osteoblastic function. *J. Bone Miner. Metab.* 25: 407-413.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 609826. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
9. Tencza, A.L., Ichikawa, S., Dang, A., Kenagy, D., McCarthy, E., Econs, M.J. and Levine, M.A. 2009. Hypophosphatemic rickets with hypercalciuria due to mutation in SLC34A3/type IIc sodium-phosphate cotransporter: presentation as hypercalciuria and nephrolithiasis. *J. Clin. Endocrinol. Metab.* 94: 4433-4438.

CHROMOSOMAL LOCATION

Genetic locus: Slc34a3 (mouse) mapping to 2 A3.

PRODUCT

NPT2c siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NPT2c shRNA Plasmid (m): sc-150057-SH and NPT2c shRNA (m) Lentiviral Particles: sc-150057-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NPT2c siRNA (m) is recommended for the inhibition of NPT2c expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NPT2c gene expression knockdown using RT-PCR Primer: NPT2c (m)-PR: sc-150057-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.