



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NUDT13 siRNA (m): sc-150103

BACKGROUND

NUDT13 (nudix (nucleoside diphosphate linked moiety X)-type motif 13) is a 352 amino acid protein belonging to the Nudix hydrolase family of pyrophosphatases. Nudix hydrolases contain a characteristic Nudix domain and are responsible for catalyzing the hydrolysis of nucleoside diphosphate derivatives. Existing as two isoforms due to alternative splicing events, NUDT13 is highly expressed in metastasis-suppressed chromosome 6 melanoma hybrids. The gene encoding NUDT13 maps to human chromosome 10, which houses over 1,200 genes and comprises nearly 4.5% of the human genome. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie-Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromic deafness, Wolman's syndrome, Cowden syndrome, multiple endocrine neoplasia type 2 and porphyria.

REFERENCES

1. Subramani, S. 1996. Convergence of model systems for peroxisome biogenesis. *Curr. Opin. Cell Biol.* 8: 513-518.
2. Alimova-Kost, M.V., Imreh, S., Buchman, V.L. and Ninkina, N.N. 1998. Assignment of phosphotriesterase-related gene (PTER) to human chromosome band 10p12 by *in situ* hybridization. *Cytogenet. Cell Genet.* 83: 16-17.
3. Berger, P., Young, P. and Suter, U. 2002. Molecular cell biology of Charcot-Marie-Tooth disease. *Neurogenetics* 4: 1-15.
4. Nonneman, D. and Rohrer, G.A. 2004. Comparative mapping of human chromosome 10 to pig chromosomes 10 and 14. *Anim. Genet.* 3: 338-343.
5. Kurochkin, I.V., Nagashima, T., Konagaya, A. and Schönbach, C. 2005. Sequence-based discovery of the human and rodent peroxisomal proteome. *Appl. Bioinformatics* 4: 93-104.
6. Kurochkin, I.V., Mizuno, Y., Konagaya, A., Sakaki, Y., Schönbach, C. and Okazaki, Y. 2007. Novel peroxisomal protease Tysnd1 processes PTS1- and PTS2-containing enzymes involved in β -oxidation of fatty acids. *EMBO J.* 26: 835-845.
7. Kraszevska, E. 2008. The plant Nudix hydrolase family. *Acta Biochim. Pol.* 55: 663-671.
8. Lin, J., Tian, B. and Hua, Y. 2008. Structural and functional diversity of Nudix fold. *Protein Pept. Lett.* 15: 108-112.
9. Garza, J.A., Ilangoan, U., Hinck, A.P. and Barnes, L.D. 2009. Kinetic, dynamic, ligand binding properties, and structural models of a dual-substrate specific nudix hydrolase from *Schizosaccharomyces pombe*. *Biochemistry* 48: 6224-6239.

CHROMOSOMAL LOCATION

Genetic locus: Nudt13 (mouse) mapping to 14 A3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

NUDT13 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NUDT13 shRNA Plasmid (m): sc-150103-SH and NUDT13 shRNA (m) Lentiviral Particles: sc-150103-V as alternate gene silencing products.

For independent verification of NUDT13 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-150103A, sc-150103B and sc-150103C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NUDT13 siRNA (m) is recommended for the inhibition of NUDT13 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NUDT13 gene expression knockdown using RT-PCR Primer: NUDT13 (m)-PR: sc-150103-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.