



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ORP-11 siRNA (m): sc-151319

BACKGROUND

Members of the oxysterol-binding protein (OSBP) family function as intracellular lipid receptors. OSBPs are involved in lipid metabolism and signal transduction, as well as vesicle transport, and can translocate to the periphery of Golgi membranes when they are bound to oxysterols. ORPs (OSBP-related proteins) belong to a subfamily of OSBPs and consists of ORP-1 through ORP-11. The ORPs have a highly conserved OSBP-type sterol-binding region and a pleckstrin homology domain. They strongly bind to phosphatidic acid and weakly bind to phosphatidylinositol 3-phosphate. The gene encoding the 747 amino acid ORP-11 (oxysterol-binding protein-related protein 11) protein maps to human chromosome 3q21.2, which spans 200 million base pairs and encodes between 1,100 and 1,500 genes.

REFERENCES

1. Jaworski, C.J., Moreira, E., Li, A., Lee, R. and Rodriguez, I.R. 2001. A family of 12 human genes containing oxysterol-binding domains. *Genomics* 78: 185-196.
2. Lehto, M., Laitinen, S., Chinetti, G., Johansson, M., Ehnholm, C., Staels, B., Ikonen, E. and Olkkonen, V.M. 2001. The OSBP-related protein family in humans. *J. Lipid Res.* 42: 1203-1213.
3. Anniss, A.M., Apostolopoulos, J., Dworkin, S., Purton, L.E. and Sparrow, R.L. 2002. An oxysterol-binding protein family identified in the mouse. *DNA Cell Biol.* 21: 571-580.
4. Lehto, M. and Olkkonen, V.M. 2003. The OSBP-related proteins: a novel protein family involved in vesicle transport, cellular lipid metabolism, and cell signalling. *Biochim. Biophys. Acta* 1631: 1-11.
5. Olkkonen, V.M. and Levine, T.P. 2004. Oxysterol binding proteins: in more than one place at one time? *Biochem. Cell Biol.* 82: 87-98.
6. Skirpan, A.L., Dowd, P.E., Sijacic, P., Jaworski, C.J., Gilroy, S. and Kao, T.H. 2006. Identification and characterization of PiORP1, a *Petunia* oxysterol-binding-protein related protein involved in receptor-kinase mediated signalling in pollen, and analysis of the ORP gene family in *Arabidopsis*. *Plant Mol. Biol.* 61: 553-565.
7. Suchanek, M., Hynynen, R., Wohlfahrt, G., Lehto, M., Johansson, M., Saarinen, H., Radzikowska, A., Thiele, C. and Olkkonen, V.M. 2007. The mammalian oxysterol-binding protein-related proteins (ORPs) bind 25-hydroxycholesterol in an evolutionarily conserved pocket. *Biochem. J.* 405: 473-480.
8. Lessmann, E., Ngo, M., Leitges, M., Minguet, S., Ridgway, N.D. and Huber, M. 2007. Oxysterol-binding protein-related protein (ORP) 9 is a PDK-2 substrate and regulates Akt phosphorylation. *Cell. Signal.* 19: 384-392.
9. Yan, D., Mäyränpää, M.I., Wong, J., Perttilä, J., Lehto, M., Jauhainen, M., Kovanen, P.T., Ehnholm, C., Brown, A.J. and Olkkonen, V.M. 2008. OSBP-related protein 8 (ORP8) suppresses ABCA1 expression and cholesterol efflux from macrophages. *J. Biol. Chem.* 283: 332-340.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Osbpl11* (mouse) mapping to 16 B3.

PRODUCT

ORP-11 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ORP-11 shRNA Plasmid (m): sc-151319-SH and ORP-11 shRNA (m) Lentiviral Particles: sc-151319-V as alternate gene silencing products.

For independent verification of ORP-11 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-151319A, sc-151319B and sc-151319C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ORP-11 siRNA (m) is recommended for the inhibition of ORP-11 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ORP-11 gene expression knockdown using RT-PCR Primer: ORP-11 (m)-PR: sc-151319-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.