



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



P504S siRNA (m): sc-151968

BACKGROUND

P504S, also known as AMACR (α -methylacyl-CoA racemase), 2-methylacyl-CoA racemase or RACE, is an enzyme belonging to the caiB/baiF CoA-transferase family. Localizing to the peroxisome and mitochondrion, P504S plays a role in the metabolism of branched-chain fatty acids and bile acid intermediates. More specifically, P504S catalyzes the conversion of pristanoyl-CoA and C27-bile acyl-CoAs to their (S)-stereoisomers which can then be degraded by peroxisomal β -oxidation. Mutations in the gene encoding P504S can lead to AMACR deficiency, a disease characterized by increased concentrations of pristanic acid that is associated with adult onset sensory motor neuropathy, and/or CBAS4 (congenital bile acid synthesis defect type 4), a disorder characterized by intrahepatic cholestasis, absence of cholic acid from bile, neonatal jaundice and bile duct deficiency. In addition, P504S is overexpressed in prostate cancer and is believed to be functionally important for prostate cancer cell growth.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604489. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Lee, W.A. 2006. α -methylacyl-CoA-racemase expression in adenocarcinoma, dysplasia and non-neoplastic epithelium of the stomach. *Oncology* 71: 246-250.
3. Chen, W., et al. 2007. Molecular cloning and preliminary analysis of the human α -methylacyl-CoA racemase promoter. *Mol. Biol. Rep.* 36: 423-430.
4. Shilo, K., et al. 2007. α -methylacyl CoA racemase in pulmonary adenocarcinoma, squamous cell carcinoma, and neuroendocrine tumors: expression and survival analysis. *Arch. Pathol. Lab. Med.* 131: 1555-1560.
5. Daugherty, S.E., et al. 2007. Polymorphic variants in α -methylacyl-CoA racemase and prostate cancer. *Prostate* 67: 1487-1497.
6. Levin, A.M., et al. 2007. Sequence variation in α -methylacyl-CoA racemase and risk of early-onset and familial prostate cancer. *Prostate* 67: 1507-1513.
7. Daugherty, S.E., et al. 2007. Variants in the α -methylacyl-CoA racemase gene and the association with advanced distal colorectal adenoma. *Cancer Epidemiol. Biomarkers Prev.* 16: 1536-1542.

CHROMOSOMAL LOCATION

Genetic locus: Amacr (mouse) mapping to 15 A1.

PRODUCT

P504S siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see P504S shRNA Plasmid (m): sc-151968-SH and P504S shRNA (m) Lentiviral Particles: sc-151968-V as alternate gene silencing products.

For independent verification of P504S (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-151968A, sc-151968B and sc-151968C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

P504S siRNA (m) is recommended for the inhibition of P504S expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

P504S (H-2): sc-514424 is recommended as a control antibody for monitoring of P504S gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor P504S gene expression knockdown using RT-PCR Primer: P504S (m)-PR: sc-151968-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.