



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



PARS2 siRNA (m): sc-152030

BACKGROUND

The fidelity of protein synthesis requires efficient discrimination of amino acid substrates by aminoacyl-tRNA synthetases. Aminoacyl-tRNA synthetases function to catalyze the aminoacylation of tRNAs by their corresponding amino acids, thus linking amino acids with tRNA-contained nucleotide triplets. PARS2 (prolyl-tRNA synthetase 2, mitochondrial), also known as MT-PRORS, is a 475 amino acid protein belonging to the class-II aminoacyl-tRNA synthetase family. Localized to the mitochondrial matrix, PARS2 utilizes ATP to catalyze the linking of L-proline and tRNA(Pro). PARS2 has been found to have no significant similarity to ProRS, its cytosolic counterpart.

REFERENCES

1. Kaiser, E., Hu, B., Becher, S., Eberhard, D., Schray, B., Baack, M., Hameister, H. and Knippers, R. 1994. The human EPRS locus (formerly the QARS locus): a gene encoding a class I and a class II aminoacyl-tRNA synthetase. *Genomics* 19: 280-290.
2. Rho, S.B., Lee, J.S., Jeong, E.J., Kim, K.S., Kim, Y.G. and Kim, S. 1998. A multifunctional repeated motif is present in human bifunctional tRNA synthetase. *J. Biol. Chem.* 273: 11267-11273.
3. Quevillon, S., Robinson, J.C., Berthonneau, E., Siatecka, M. and Mirande, M. 1999. Macromolecular assemblage of aminoacyl-tRNA synthetases: identification of protein-protein interactions and characterization of a core protein. *J. Mol. Biol.* 285: 183-195.
4. Jeong, E.J., Hwang, G.S., Kim, K.H., Kim, M.J., Kim, S. and Kim, K.S. 2000. Structural analysis of multifunctional peptide motifs in human bifunctional tRNA synthetase: identification of RNA-binding residues and functional implications for tandem repeats. *Biochemistry* 39: 15775-15782.
5. Kang, J., Kim, T., Ko, Y.G., Rho, S.B., Park, S.G., Kim, M.J., Kwon, H.J. and Kim, S. 2000. Heat shock protein 90 mediates protein-protein interactions between human aminoacyl-tRNA synthetases. *J. Biol. Chem.* 275: 31682-31688.
6. Sang Lee, J., Gyu Park, S., Park, H., Seol, W., Lee, S. and Kim, S. 2002. Interaction network of human aminoacyl-tRNA synthetases and subunits of elongation factor 1 complex. *Biochem. Biophys. Res. Commun.* 291: 158-164.
7. Bonnefond, L., Fender, A., Rudinger-Thirion, J., Giege, R., Florentz, C. and Sissler, M. 2005. Toward the full set of human mitochondrial aminoacyl-tRNA synthetases: characterization of AspRS and TyrRS. *Biochemistry* 44: 4805-4816.
8. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612036. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Pars2 (mouse) mapping to 4 C7.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

PARS2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PARS2 shRNA Plasmid (m): sc-152030-SH and PARS2 shRNA (m) Lentiviral Particles: sc-152030-V as alternate gene silencing products.

For independent verification of PARS2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152030A, sc-152030B and sc-152030C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PARS2 siRNA (m) is recommended for the inhibition of PARS2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PARS2 gene expression knockdown using RT-PCR Primer: PARS2 (m)-PR: sc-152030-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.