



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PHI-1 siRNA (m): sc-152222

BACKGROUND

PHI-1, also known as PPP1R14B (protein phosphatase 1, regulatory (inhibitor) subunit 14B), PLCB3N (phospholipase C- β -3 neighbouring gene protein), PNG or SOM172, is a 147 amino acid coiled-coil protein that belongs to the PP1 inhibitor family. Ubiquitously expressed at low levels, PHI-1 localizes to cytoplasm and functions as an inhibitor of PP1. PHI-1 blocks myosin light chain dephosphorylation and interacts with MYPT1, an important regulator of cell migration, adhesion and retraction that may function as a tumor suppressor by regulating Rho-dependent amoeboid cell behavior in metastasis. Phosphorylated primarily on threonine 57 and an unknown serine by PKC, PHI-1 exhibits a 50-fold increase in inhibitory activity during phosphorylation. The gene that encodes PHI-1 maps to human chromosome 11q13.1.

REFERENCES

1. Lagercrantz, J., Kedra, D., Carson, E., Nordenskjöld, M., Dumanski, J.P., Weber, G. and Piehl, F. 1996. Sequence and expression of the mouse homologue to human phospholipase C β 3 neighboring gene. *Biochem. Biophys. Res. Commun.* 223: 335-340.
2. Lagercrantz, J., Carson, E., Larsson, C., Nordenskjöld, M. and Weber, G. 1996. Isolation and characterization of a novel gene close to the human phosphoinositide-specific phospholipase C β 3 gene on chromosomal region 11q13. *Genomics* 31: 380-384.
3. Ceulemans, H., Stalmans, W. and Bollen, M. 2002. Regulator-driven functional diversification of protein phosphatase-1 in eukaryotic evolution. *Bioessays* 24: 371-381.
4. Cerutti, J.M., Delcelo, R., Amadei, M.J., Nakabashi, C., Maciel, R.M., Peterson, B., Shoemaker, J. and Riggins, G.J. 2004. A preoperative diagnostic test that distinguishes benign from malignant thyroid carcinoma based on gene expression. *J. Clin. Invest.* 113: 1234-1242.
5. Liu, Q.R., Gong, J.P. and Uhl, G.R. 2005. Families of protein phosphatase 1 modulators activated by protein kinases A and C: focus on brain. *Prog. Nucleic Acid Res. Mol. Biol.* 79: 371-404.
6. Weiser, D.C., Row, R.H. and Kimelman, D. 2009. Rho-regulated myosin phosphatase establishes the level of protrusive activity required for cell movements during zebrafish gastrulation. *Development* 136: 2375-2384.
7. Drgonova, J., Zimonjic, D.B., Hall, F.S. and Uhl, G.R. 2010. Effect of KEPI (Ppp1r14c) deletion on morphine analgesia and tolerance in mice of different genetic backgrounds: when a knockout is near a relevant quantitative trait locus. *Neuroscience* 165: 882-895.

CHROMOSOMAL LOCATION

Genetic locus: Ppp1r14b (mouse) mapping to 19 A.

PRODUCT

PHI-1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PHI-1 shRNA Plasmid (m): sc-152222-SH and PHI-1 shRNA (m) Lentiviral Particles: sc-152222-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PHI-1 siRNA (m) is recommended for the inhibition of PHI-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

PHI-1 (D-8): sc-514759 is recommended as a control antibody for monitoring of PHI-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PHI-1 gene expression knockdown using RT-PCR Primer: PHI-1 (m)-PR: sc-152222-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.