



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PHKG1 siRNA (m): sc-152225

BACKGROUND

PHKG1 (phosphorylase kinase subunit γ 1), is a subunit of phosphorylase kinase (PHK) that belongs to the Ser/Thr protein kinase family. PHK is a hexadecameric protein composed of four α chains, four β chains, four γ chains and four δ chains. The γ chains are catalytic chains, the α and β chains are regulatory chains and the δ chains are calmodulins. PHKG1 contains two calmodulin-binding domains and one protein kinase domain. As the catalytic chain of PHK, PHKG1 is responsible for catalyzing the phosphorylation and activation of glycogen phosphorylase and therefore it plays an important role in the glycogenolytic pathway. Mutations in the gene encoding PHKG1 can lead to PHK deficiency and result in glycogen storage disease type 9C (GSD9C), also known as autosomal liver glycogenosis.

REFERENCES

- Hanks, S.K. 1989. Messenger ribonucleic acid encoding an apparent isoform of phosphorylase kinase catalytic subunit is abundant in the adult testis. *Mol. Endocrinol.* 3: 110-116.
- Calalb, M.B., Fox, D.T. and Hanks, S.K. 1992. Molecular cloning and enzymatic analysis of the rat homolog of "PhK- γ T", an isoform of phosphorylase kinase catalytic subunit. *J. Biol. Chem.* 267: 1455-1463.
- Liu, L., Rannels, S.R., Falconieri, M., Phillips, K.S., Wolpert, E.B. and Weaver, T.E. 1996. The testis isoform of the phosphorylase kinase catalytic subunit (PhK- γ T) plays a critical role in regulation of glycogen mobilization in developing lung. *J. Biol. Chem.* 271: 11761-11766.
- Maichele, A.J., Burwinkel, B., Maire, I., Sovik, O. and Kilimann, M.W. 1996. Mutations in the testis/liver isoform of the phosphorylase kinase γ subunit (PHKG2) cause autosomal liver glycogenosis in the gsd rat and in humans. *Nat. Genet.* 14: 337-340.
- Burwinkel, B., Shiomi, S., Al Zaben, A. and Kilimann, M.W. 1998. Liver glycogenosis due to phosphorylase kinase deficiency: PHKG2 gene structure and mutations associated with cirrhosis. *Hum. Mol. Genet.* 7: 149-154.
- Burwinkel, B., Tanner, M.S. and Kilimann, M.W. 2000. Phosphorylase kinase deficient liver glycogenosis: progression to cirrhosis in infancy associated with PHKG2 mutations (H14.and L225R). *J. Med. Genet.* 37: 376-377.
- Burwinkel, B., Rootwelt, T., Kvittingen, E.A., Chakraborty, P.K. and Kilimann, M.W. 2003. Severe phenotype of phosphorylase kinase-deficient liver glycogenosis with mutations in the PHKG2 gene. *Pediatr. Res.* 54: 834-839.
- Chen, C.S., Chen, N.J., Lin, L.W., Hsieh, C.C., Chen, G.W. and Hsieh, M.T. 2006. Effects of Scutellariae Radix on gene expression in HEK 293 cells using cDNA microarray. *J. Ethnopharmacol.* 105: 346-351.
- Beauchamp, N.J., Dalton, A., Ramaswami, U., Niinikoski, H., Mention, K., Kenny, P., Kolho, K.L., Raiman, J., Walter, J., Treacy, E., Tanner, S. and Sharrard, M. 2007. Glycogen storage disease type IX: High variability in clinical phenotype. *Mol. Genet. Metab.* 92: 88-99.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Phkg1 (mouse) mapping to 5 G1.3.

PRODUCT

PHKG1 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PHKG1 shRNA Plasmid (m): sc-152225-SH and PHKG1 shRNA (m) Lentiviral Particles: sc-152225-V as alternate gene silencing products.

For independent verification of PHKG1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152225A and sc-152225B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PHKG1 siRNA (m) is recommended for the inhibition of PHKG1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PHKG1 gene expression knockdown using RT-PCR Primer: PHKG1 (m)-PR: sc-152225-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.