



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PPC-DC siRNA (m): sc-152407



The Power to Question

BACKGROUND

Phosphopantethenoylcysteine decarboxylase (PPC-DC) is a 204 amino acid protein that plays a role in the biosynthesis of coenzyme A (CoA) from pantothenate (vitamin B). CoA is an essential cofactor in all living organisms and is involved in several key biochemical pathways, including the tricarboxylic acid cycle and fatty acid metabolism. Altered CoA levels are associated with aberrant mitosis and meiosis in flies and the neurodegenerative Hallervorden-Spatz syndrome in humans. The biosynthesis of CoA from pantothenate requires several steps: the phosphorylation of pantothenate, the conversion of 4'-phosphopantethenate to 4'-phosphopantetheine, the adenylation by phosphopantetheine adenyltransferase to form dephospho-CoA and the phosphorylation by dephospho-CoA kinase to form CoA. PPC-DC plays a direct role in this pathway by converting 4'-phosphopantethenate into 4'-phosphopantetheine. Potentially forming a homotrimer, PPC-DC has two named isoforms produced by alternative splicing.

REFERENCES

- Strauss, E., Kinsland, C., Ge, Y., McLafferty, F.W. and Begley, T.P. 2001. Phosphopantethenoylcysteine synthetase from *Escherichia coli*. Identification and characterization of the last unidentified coenzyme A biosynthetic enzyme in bacteria. *J. Biol. Chem.* 276: 13513-13516.
- Daugherty, M., Polanuyer, B., Farrell, M., Scholle, M., Lykidis, A., de Crecy-Lagard, V. and Osterman, A. 2002. Complete reconstitution of the human coenzyme A biosynthetic pathway via comparative genomics. *J. Biol. Chem.* 277: 21431-21439.
- Zhyvoloup, A., Nemazanyy, I., Babich, A., Panasyuk, G., Pobigailo, N., Vudmaska, M., Naidenov, V., Kukharenko, O., Palchevskii, S., Savinska, L., Ovcharenko, G., Verdier, F., Valovka, T., Fenton, T., Rebholz, H., et al. 2002. Molecular cloning of CoA Synthase. The missing link in CoA biosynthesis. *J. Biol. Chem.* 277: 22107-22110.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 609854. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Kupke, T., Hernández-Acosta, P. and Culianez-Macià, F.A. 2003. 4'-phosphopantetheine and coenzyme A biosynthesis in plants. *J. Biol. Chem.* 278: 38229-38237.
- Zhyvoloup, A., Nemazanyy, I., Panasyuk, G., Valovka, T., Fenton, T., Rebholz, H., Wang, M.L., Foxon, R., Lyzogubov, V., Usenko, V., Kyyamova, R., Gorbenko, O., Matsuka, G., Filonenko, V. and Gout, I.T. 2003. Subcellular localization and regulation of coenzyme A synthase. *J. Biol. Chem.* 278: 50316-50321.
- Strauss, E., Zhai, H., Brand, L.A., McLafferty, F.W. and Begley, T.P. 2004. Mechanistic studies on phosphopantethenoylcysteine decarboxylase: trapping of an enethiolate intermediate with a mechanism-based inactivating agent. *Biochemistry* 43: 15520-15533.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Ppcdc* (mouse) mapping to 9 B.

PRODUCT

PPC-DC siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PPC-DC shRNA Plasmid (m): sc-152407-SH and PPC-DC shRNA (m) Lentiviral Particles: sc-152407-V as alternate gene silencing products.

For independent verification of PPC-DC (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-152407A, sc-152407B and sc-152407C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PPC-DC siRNA (m) is recommended for the inhibition of PPC-DC expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PPC-DC gene expression knockdown using RT-PCR Primer: PPC-DC (m)-PR: sc-152407-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.