



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TAAR8A siRNA (m): sc-154038

BACKGROUND

Trace amines are biogenic amines present in very low levels in mammalian tissues. Although some trace amines have clearly defined roles as neurotransmitters in invertebrates, the extent to which they function as true neurotransmitters in vertebrates is unclear. Trace amines are likely involved in a variety of physiological functions that have yet to be fully understood. TAAR8A (trace amine-associated receptor 8a), also known as Tar-8a or Gm230, is a 344 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor 1 family. The gene that encodes TAAR8A maps to mouse chromosome 10 A4.

REFERENCES

1. Borowsky, B., Adham, N., Jones, K.A., Raddatz, R., Artymyshyn, R., Ogozalek, K.L., Durkin, M.M., Lakhani, P.P., Bonini, J.A., Pathirana, S., Boyle, N., Pu, X., Kouranova, E., Lichtblau, H., Ochoa, F.Y., Brancheck, T.A., et al. 2001. Trace amines: identification of a family of mammalian G protein-coupled receptors. Proc. Natl. Acad. Sci. USA 98: 8966-8971.
2. Lindemann, L., Ebeling, M., Kratochwil, N.A., Bunzow, J.R., Grandy, D.K. and Hoener, M.C. 2005. Trace amine-associated receptors form structurally and functionally distinct subfamilies of novel G protein-coupled receptors. Genomics 85: 372-385.
3. Zucchi, R., Chiellini, G., Scanlan, T.S. and Grandy, D.K. 2006. Trace amine-associated receptors and their ligands. Br. J. Pharmacol. 149: 967-978.
4. Liberles, S.D. and Buck, L.B. 2006. A second class of chemosensory receptors in the olfactory epithelium. Nature 442: 645-650.
5. Fleischer, J., Schwarzenbacher, K. and Breer, H. 2007. Expression of trace amine-associated receptors in the Grueneberg ganglion. Chem. Senses 32: 623-631.
6. Liberles, S.D. 2009. Trace amine-associated receptors are olfactory receptors in vertebrates. Ann. N.Y. Acad. Sci. 1170: 168-172.
7. Sotnikova, T.D., Caron, M.G. and Gainetdinov, R.R. 2009. Trace amine-associated receptors as emerging therapeutic targets. Mol. Pharmacol. 76: 229-235.
8. Chiellini, G., Erba, P., Carnicelli, V., Manfredi, C., Frascarelli, S., Ghelardoni, S., Mariani, G. and Zucchi, R. 2012. Distribution of exogenous [¹²⁵I]-3-iodothyronamine in mouse *in vivo*: relationship with trace amine-associated receptors. J. Endocrinol. 213: 223-230.

CHROMOSOMAL LOCATION

Genetic locus: Taar8a (mouse) mapping to 10 A4.

PRODUCT

TAAR8A siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TAAR8A shRNA Plasmid (m): sc-154038-SH and TAAR8A shRNA (m) Lentiviral Particles: sc-154038-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TAAR8A siRNA (m) is recommended for the inhibition of TAAR8A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TAAR8A gene expression knockdown using RT-PCR Primer: TAAR8A (m)-PR: sc-154038-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.