



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TAPBPL siRNA (m): sc-154069

BACKGROUND

TAPBPL (TAP binding protein-like), also known as TAPBPR or TAPBP-R, is a 468 amino acid protein that contains one Ig-like (immunoglobulin-like) C1-type domain and one Ig-like V-type domain. Localized to the membrane of both the endoplasmic reticulum (ER) and the microsome, TAPBPL is a single-pass type I membrane protein that is similar to TPN (also known as TAPBP or Tapasin), a transmembrane glycoprotein that belongs to the variable-constant Ig superfamily. TPN functions to link the ER-associated antigen transporter TAP with major histocompatibility complex (MHC) class I molecules, thereby mediating peptide loading onto MHC proteins. Due to its similarity with TPN, TAPBPL is thought to play a role in antigen processing events within the ER.

REFERENCES

- Sadasivan, B., Lehner, P.J., Ortmann, B., Spies, T. and Cresswell, P. 1996. Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* 5: 103-114.
- Ortmann, B., Copeman, J., Lehner, P.J., Sadasivan, B., Herberg, J.A., Grandea, A.G., Riddell, S.R., Tampe, R., Spies, T., Trowsdale, J. and Cresswell, P. 1997. A critical role for tapasin in the assembly and function of multimeric MHC class I-TAP complexes. *Science* 277: 1306-1309.
- Herberg, J.A., Sgouros, J., Jones, T., Copeman, J., Humphray, S.J., Sheer, D., Cresswell, P., Beck, S. and Trowsdale, J. 1998. Genomic analysis of the Tapasin gene, located close to the TAP loci in the MHC. *Eur. J. Immunol.* 28: 459-467.
- Mayer, W.E. and Klein, J. 2001. Is tapasin a modified Mhc class I molecule? *Immunogenetics* 53: 719-723.
- Teng, M.S., Stephens, R., Du Pasquier, L., Freeman, T., Lindquist, J.A. and Trowsdale, J. 2002. A human TAPBP (TAPASIN)-related gene, TAPBP-R. *Eur. J. Immunol.* 32: 1059-1068.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607081. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Rizvi, S.M. and Raghavan, M. 2006. Direct peptide-regulatable interactions between MHC class I molecules and tapasin. *Proc. Natl. Acad. Sci. USA* 103: 18220-18225.
- Chambers, J.E., Jessop, C.E. and Bulleid, N.J. 2008. Formation of a major histocompatibility complex class I tapasin disulfide indicates a change in spatial organization of the peptide-loading complex during assembly. *J. Biol. Chem.* 283: 1862-1869.

CHROMOSOMAL LOCATION

Genetic locus: *Tapbp1* (mouse) mapping to 6 F3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TAPBPL siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TAPBPL shRNA Plasmid (m): sc-154069-SH and TAPBPL shRNA (m) Lentiviral Particles: sc-154069-V as alternate gene silencing products.

For independent verification of TAPBPL (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154069A, sc-154069B and sc-154069C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TAPBPL siRNA (m) is recommended for the inhibition of TAPBPL expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TAPBPL gene expression knockdown using RT-PCR Primer: TAPBPL (m)-PR: sc-154069-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.