



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Tctex1D2 siRNA (m): sc-154151

BACKGROUND

Dyneins are multi-subunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Cytoplasmic or axonemal dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; the complex transports cellular cargos towards the central region of the cell. Tctex1, also designated CW-1 or TCTEL1, is expressed in heart, placenta, skeletal muscle, kidney, pancreas, spleen, prostate, testis, ovary, ileum and colon, where it functions as a cytoplasmic dynein light chain and is found in a complex with Na⁺ CP type Xα. Tctex1D2 (Tctex1 domain containing 2) is a 144 amino acid protein that belongs to the dynein light chain Tctex-type family and may function in a similar manner to Tctex1.

REFERENCES

- Harrison, A., Olds-Clarke, P. and King, S.M. 1998. Identification of the t complex-encoded cytoplasmic dynein light chain Tctex1 in inner arm I1 supports the involvement of flagellar dyneins in meiotic drive. *J. Cell Biol.* 140: 1137-1147.
- Harrison, A. and King, S.M. 2000. The molecular anatomy of dynein. *Essays Biochem.* 35: 75-87.
- Wilson, M.J., Salata, M.W., Susalka, S.J. and Pfister, K.K. 2001. Light chains of mammalian cytoplasmic dynein: identification and characterization of a family of LC8 light chains. *Cell Motil. Cytoskeleton* 49: 229-240.
- DiBella, L.M., Benashski, S.E., Tedford, H.W., Harrison, A., Patel-King, R.S. and King, S.M. 2001. The Tctex1/Tctex2 class of dynein light chains. Dimerization, differential expression, and interaction with the LC8 protein family. *J. Biol. Chem.* 276: 14366-14373.
- Weise, A., Starke, H., Mrasek, K., Claussen, U. and Liehr, T. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
- DiBella, L.M., Gorbatyuk, O., Sakato, M., Wakabayashi, K., Patel-King, R.S., Pazour, G.J., Witman, G.B. and King, S.M. 2005. Differential light chain assembly influences outer arm dynein motor function. *Mol. Biol. Cell* 16: 5661-5674.
- Williams, J.C., Roulhac, P.L., Roy, A.G., Vallee, R.B., Fitzgerald, M.C. and Hendrickson, W.A. 2007. Structural and thermodynamic characterization of a cytoplasmic dynein light chain-intermediate chain complex. *Proc. Natl. Acad. Sci. USA* 104: 10028-10033.
- Githui, E.K., De Villiers, E.P. and McArthur, A.G. 2009. Plasmodium possesses dynein light chain classes that are unique and conserved across species. *Infect. Genet. Evol.* 9: 337-343.
- Palmer, K.J., Hughes, H. and Stephens, D.J. 2009. Specificity of cytoplasmic dynein subunits in discrete membrane-trafficking steps. *Mol. Biol. Cell* 20: 2885-2899.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Tctex1d2 (mouse) mapping to 16 B3.

PRODUCT

Tctex1D2 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Tctex1D2 shRNA Plasmid (m): sc-154151-SH and Tctex1D2 shRNA (m) Lentiviral Particles: sc-154151-V as alternate gene silencing products.

For independent verification of Tctex1D2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154151A and sc-154151B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Tctex1D2 siRNA (m) is recommended for the inhibition of Tctex1D2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Tctex1D2 gene expression knockdown using RT-PCR Primer: Tctex1D2 (m)-PR: sc-154151-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.