



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TFB2M siRNA (m): sc-154229

BACKGROUND

TFB2M (transcription factor B2, mitochondrial), also known as NS5ATP5, Hkp1, Mitochondrial 12S rRNA dimethylase 2, Hepatitis C virus NS5A-transactivated protein 5 or mitochondrial Dimethyladenosine transferase 2, is a 396 amino acid mitochondrial protein that belongs to the rRNA adenine N⁶-methyltransferase family. Expressed ubiquitously, TFB2M is an S-adenosyl-L-methionine-dependent methyltransferase which specifically dimethylates mitochondrial 12S rRNA at the conserved stem loop. TFB2M is required for basal transcription of mitochondrial DNA and stimulates transcription independently of its methyltransferase activity. Compared to TFB1M, TFB2M has less methyltransferase activity but activates transcription of mitochondrial DNA more efficiently.

REFERENCES

- Falkenberg, M., Gaspari, M., Rantanen, A., Trifunovic, A., Larsson, N.G. and Gustafsson, C.M. 2002. Mitochondrial transcription factors B1 and B2 activate transcription of human mtDNA. *Nat. Genet.* 31: 289-294.
- Gleyzer, N., Vercauteren, K. and Scarpulla, R.C. 2005. Control of mitochondrial transcription specificity factors (TFB1M and TFB2M) by nuclear respiratory factors (NRF-1 and NRF-2) and PGC-1 family coactivators. *Mol. Cell. Biol.* 25: 1354-1366.
- Dubessay, P., Garreau-Balandier, I., Jarrousse, A.S., Fleuriet, A., Sion, B., Debise, R. and Alzari, S. 2007. Aging impact on biochemical activities and gene expression of *Drosophila melanogaster* mitochondria. *Biochimie* 89: 988-1001.
- Cotney, J., Wang, Z. and Shadel, G.S. 2007. Relative abundance of the human mitochondrial transcription system and distinct roles for h-mtTFB1 and h-mtTFB2 in mitochondrial biogenesis and gene expression. *Nucleic Acids Res.* 35: 4042-4054.
- Alonso-Montes, C., Castro, M.G., Reguero, J.R., Perrot, A., Ozcelik, C., Geier, C., Posch, M.G., Moris, C., Alvarez, V., Ruiz-Ortega, M. and Coto, E. 2008. Mitochondrial transcription factors TFA, TFB1 and TFB2: a search for DNA variants/haplotypes and the risk of cardiac hypertrophy. *Dis. Markers* 25: 131-139.
- Adán, C., Matsushima, Y., Hernández-Sierra, R., Marco-Ferreres, R., Fernández-Moreno, M.A., González-Vioque, E., Calleja, M., Aragón, J.J., Kaguni, L.S. and Garesse, R. 2008. Mitochondrial transcription factor B2 is essential for metabolic function in *Drosophila melanogaster* development. *J. Biol. Chem.* 283: 12333-12342.
- Alvarez, V., Corao, A.I., Sánchez-Ferrero, E., De Mena, L., Alonso-Montes, C., Huerta, C., Blázquez, M., Ribacoba, R., Guisasola, L.M., Salvador, C., García-Castro, M. and Coto, E. 2008. Mitochondrial transcription factor A (TFAM) gene variation in Parkinson's disease. *Neurosci. Lett.* 432: 79-82.
- Sánchez-Ferrero, E., Coto, E., Blázquez, M., Ribacoba, R., Guisasola, L.M., Salvador, C. and Alvarez, V. 2008. Mutational screening of the Mitochondrial transcription factors B1 and B2 (TFB1M and TFB2M) in Parkinson's disease. *Parkinsonism Relat. Disord.* 15: 468-470.

CHROMOSOMAL LOCATION

Genetic locus: Tfb2m (mouse) mapping to 1 H4.

PRODUCT

TFB2M siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TFB2M shRNA Plasmid (m): sc-154229-SH and TFB2M shRNA (m) Lentiviral Particles: sc-154229-V as alternate gene silencing products.

For independent verification of TFB2M (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154229A, sc-154229B and sc-154229C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TFB2M siRNA (m) is recommended for the inhibition of TFB2M expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TFB2M gene expression knockdown using RT-PCR Primer: TFB2M (m)-PR: sc-154229-PR (20 µl). Annealing temperature for the primers should be 55-60°C and the extension temperature should be 68-72°C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.