



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TIN-Ag siRNA (m): sc-154275

BACKGROUND

TIN-Ag (Tubulointerstitial nephritis antigen), also known as TIN1 or TIN2, is a 476 amino acid secreted protein that contains one SMB (somatomedin-B) domain and localizes to the extracellular matrix. Expressed during development in corneal tissue, kidney cortex and small intestine, TIN-Ag is a basement membrane glycoprotein that interacts with Integrin $\alpha 3$ and Integrin $\alpha 5$ and, via this interaction, mediates the adhesion of proximal tubule epithelial cells. Additionally, TIN-Ag is thought to play a role in the regulation of telomere length, possibly stabilizing the TRF2 (telomeric repeat binding factor 2) complex that is responsible for telomere elongation. Antibodies against TIN-Ag are present in the sera of tubulointerstitial nephritis-affected patients, suggesting a role for TIN-Ag in the pathogenesis of tubulointerstitial nephritis. Two isoforms of TIN-Ag are expressed due to alternative splicing events.

REFERENCES

- Katz, A., Fish, A.J., Santamaria, P., Nevins, T.E., Kim, Y. and Butkowski, R.J. 1992. Role of antibodies to tubulointerstitial nephritis antigen in human anti-tubular basement membrane nephritis associated with membranous nephropathy. *Am. J. Med.* 93: 691-698.
- Chen, Y., Krishnamurti, U., Wayner, E.A., Michael, A.F. and Charonis, A.S. 1996. Receptors in proximal tubular epithelial cells for tubulointerstitial nephritis antigen. *Kidney Int.* 49: 153-157.
- Nelson, T.R., Kim, Y., Michael, A.F., Butkowski, R.J. and Charonis, A.S. 1998. Tubulointerstitial nephritis antigen (TIN-Ag) is expressed in distinct segments of the developing human nephron. *Connect. Tissue Res.* 37: 53-60.
- Kim, S.H., Kaminker, P. and Campisi, J. 1999. TIN2, a new regulator of telomere length in human cells. *Nat. Genet.* 23: 405-412.
- Kanwar, Y.S., Kumar, A., Yang, Q., Tian, Y., Wada, J., Kashihara, N. and Wallner, E.I. 1999. Tubulointerstitial nephritis antigen: an extracellular matrix protein that selectively regulates tubulogenesis vs. glomerulogenesis during mammalian renal development. *Proc. Natl. Acad. Sci. USA* 96: 11323-11328.
- Ikeda, M., Takemura, T., Hino, S. and Yoshioka, K. 2000. Molecular cloning, expression, and chromosomal localization of a human tubulointerstitial nephritis antigen. *Biochem. Biophys. Res. Commun.* 268: 225-230.
- Zhou, B., Nelson, T.R., Kashtan, C., Gleason, B., Michael, A.F., Vlassi, M. and Charonis, A.S. 2000. Identification of two alternatively spliced forms of human tubulointerstitial nephritis antigen (TIN-Ag). *J. Am. Soc. Nephrol.* 11: 658-668.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606749. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Ye, J.Z., Donigian, J.R., van Overbeek, M., Loayza, D., Luo, Y., Krutchinsky, A.N., Chait, B.T. and de Lange, T. 2004. TIN2 binds TRF1 and TRF2 simultaneously and stabilizes the TRF2 complex on telomeres. *J. Biol. Chem.* 279: 47264-47271.

CHROMOSOMAL LOCATION

Genetic locus: Tinag (mouse) mapping to 9 D.

PRODUCT

TIN-Ag siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TIN-Ag shRNA Plasmid (m): sc-154275-SH and TIN-Ag shRNA (m) Lentiviral Particles: sc-154275-V as alternate gene silencing products.

For independent verification of TIN-Ag (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154275A, sc-154275B and sc-154275C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TIN-Ag siRNA (m) is recommended for the inhibition of TIN-Ag expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TIN-Ag gene expression knockdown using RT-PCR Primer: TIN-Ag (m)-PR: sc-154275-PR (20 μ l). Annealing temperature for the primers should be 55-60°C and the extension temperature should be 68-72°C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.