



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TM7SF2 siRNA (m): sc-154308

BACKGROUND

Transmembrane 7 superfamily member 2 (TM7SF2, Sterol C14-reductase, 3 β -hydroxysterol Δ -reductase) is a 418 amino acid gene product that belongs to the ERG4/ERG24 family. TM7SF2 is a seven pass transmembrane protein that can localize to the membrane of the endoplasmic reticulum. TM7SF2 is involved in the conversion of lanosterol to cholesterol and, specifically, catalyzes the NADPH dependant reduction of 4,4-dimethyl-5- α -cholesta-8,14,24-trien-3- β -ol to 4,4-dimethyl-5- α -cholesta-8,24-dien-3- β -ol and NADP $^+$.

REFERENCES

1. Lemmens, I.H., Kas, K., Merregaert, J. and Van de Ven, W.J. 1998. Identification and molecular characterization of TM7SF2 in the FAUNA gene cluster on human chromosome 11q13. *Genomics* 49: 437-442.
2. Holmer, L., Pezman, A. and Worman, H.J. 1998. The human Lamin B receptor/sterol reductase multigene family. *Genomics* 54: 469-476.
3. Waterham, H.R., Koster, J., Mooyer, P., Noort Gv, G., Kelley, R.I., Wilcox, W.R., Wanders, R.J., Hennekam, R.C. and Oosterwijk, J.C. 2003. Autosomal recessive HEM/Greenberg skeletal dysplasia is caused by 3 β -hydroxysterol Δ 14-reductase deficiency due to mutations in the Lamin B receptor gene. *Am. J. Hum. Genet.* 72: 1013-1017.
4. Bennati, A.M., Castelli, M., Della Fazia, M.A., Beccari, T., Caruso, D., Servillo, G. and Roberti, R. 2006. Sterol dependent regulation of human TM7SF2 gene expression: role of the encoded 3 β -hydroxysterol Δ 14-reductase in human cholesterol biosynthesis. *Biochim. Biophys. Acta* 1761: 677-685.
5. Bennati, A.M., Schiavoni, G., Franken, S., Piobbico, D., Della Fazia, M.A., Caruso, D., De Fabiani, E., Benedetti, L., Cusella De Angelis, M.G., Gieselmann, V., Servillo, G., Beccari, T. and Roberti, R. 2008. Disruption of the gene encoding 3 β -hydroxysterol Δ -reductase (Tm7sf2) in mice does not impair cholesterol biosynthesis. *FEBS J.* 275: 5034-5047.
6. Liu, Z., Rudd, M.D., Hernandez-Gonzalez, I., Gonzalez-Robayna, I., Fan, H.Y., Zeleznik, A.J. and Richards, J.S. 2009. FSH and FOXO1 regulate genes in the sterol/steroid and lipid biosynthetic pathways in granulosa cells. *Mol. Endocrinol.* 23: 649-661.
7. Schiavoni, G., Bennati, A.M., Castelli, M., Fazia, M.A., Beccari, T., Servillo, G. and Roberti, R. 2010. Activation of TM7SF2 promoter by SREBP-2 depends on a new sterol regulatory element, a GC-box, and an inverted CCAAT-box. *Biochim. Biophys. Acta* 1801: 587-592.
8. Zwerger, M., Kolb, T., Richter, K., Karakesisoglu and I., Herrmann, H. 2010. Induction of a massive endoplasmic reticulum and perinuclear space expansion by expression of lamin B receptor mutants and the related sterol reductases TM7SF2 and DHCR7. *Mol. Biol. Cell* 21: 354-368.

CHROMOSOMAL LOCATION

Genetic locus: Tm7sf2 (mouse) mapping to 19 A.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TM7SF2 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TM7SF2 shRNA Plasmid (m): sc-154308-SH and TM7SF2 shRNA (m) Lentiviral Particles: sc-154308-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TM7SF2 siRNA (m) is recommended for the inhibition of TM7SF2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TM7SF2 gene expression knockdown using RT-PCR Primer: TM7SF2 (m)-PR: sc-154308-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.