



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



ELP6 siRNA (m): sc-154341

BACKGROUND

In *Saccharomyces cerevisiae*, the hyperphosphorylated form of RNA polymerase II (RNAP II) mediates transcription elongation, and associates with the elongator complex, which contains six subunits. The elongator complex can be separated into two subcomplexes; one consisting of Elp1, Elp2 and Elp3, and the other consisting of Elp4, Elp5 and Elp6. The elongator complex acetylates both core histones and nucleosomal substrates, and directs its activity specifically towards the N-terminal tails of Histone H3 and Histone H4. An analogous complex exists in mammals and contains a variety of proteins that are functional homologs of their yeast counterparts. ELP6 (elongation protein 6), also known as C3orf75 or TMEM103, is a 266 amino acid protein and is a component of the elongator complex. Existing as two alternatively spliced isoforms, ELP6 is involved in transcriptional regulation and may play a role in cell migration.

REFERENCES

- Winkler, G.S., Petrakis, T.G., Ethelberg, S., Tokunaga, M., Erdjument-Bromage, H., Tempst, P. and Svejstrup, J.O. 2001. RNA polymerase II elongator holoenzyme is composed of two discrete subcomplexes. *J. Biol. Chem.* 276: 32743-32749.
- Kim, J.H., Lane, W.S. and Reinberg, D. 2002. Human Elongator facilitates RNA polymerase II transcription through chromatin. *Proc. Natl. Acad. Sci. USA* 99: 1241-1246.
- Li, F., Han, Q.J., Luo, D.H., Lu, J. and Huang, B.Q. 2004. The Elp4 subunit of human Elongator complex partially complements the growth defects of yeast ELP4 deletion strain. *Yi Chuan Xue Bao* 31: 668-674.
- Nelissen, H., Fleury, D., Bruno, L., Robles, P., De Veylder, L., Traas, J., Micol, J.L., Van Montagu, M., Inze, D. and Van Lijsebettens, M. 2005. The elongata mutants identify a functional Elongator complex in plants with a role in cell proliferation during organ growth. *Proc. Natl. Acad. Sci. USA* 102: 7754-7759.
- Srokowski, E.M. and Woodhouse, K.A. 2008. Development and characterisation of novel cross-linked bio-elastomeric materials. *J. Biomater. Sci. Polym. Ed.* 19: 785-799.
- Strug, L.J., Clarke, T., Chiang, T., Chien, M., Baskurt, Z., Li, W., Dorfman, R., Bali, B., Wirrell, E., Kugler, S.L., Mandelbaum, D.E., Wolf, S.M., McGoldrick, P., Hardison, H., Novotny, E.J., Ju, J., Greenberg, D.A., Russo, J.J., et al. 2009. Centrotemporal sharp wave EEG trait in rolandic epilepsy maps to Elongator Protein Complex 4 (ELP4). *Eur. J. Hum. Genet.* 17: 1171-1181.
- Close, P., Gillard, M., Ladang, A., Jiang, Z., Papuga, J., Hawkes, N., Nguyen, L., Chapelle, J.P., Bouillenne, F., Svejstrup, J., Fillet, M. and Chariot, A. 2012. DERP6 (ELP5) and C3ORF75 (ELP6) regulate tumorigenicity and migration of melanoma cells as subunits of Elongator. *J. Biol. Chem.* 287: 32535-32545.
- Wagner, L.A., Wang, S., Wayner, E.A., Christensen, C., Perkins, S.J., Ward, G.W., Weiss, R.B., Dunn, D.M., Redd, M.J., Spangrude, G.J. and Gleich, G.J. 2013. Developing and mature human granulocytes express ELP 6 in the cytoplasm. *Hum. Antibodies* 22: 21-29.

CHROMOSOMAL LOCATION

Genetic locus: Elp6 (mouse) mapping to 9 F2.

PRODUCT

ELP6 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ELP6 shRNA Plasmid (m): sc-154341-SH and ELP6 shRNA (m) Lentiviral Particles: sc-154341-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ELP6 siRNA (m) is recommended for the inhibition of ELP6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ELP6 gene expression knockdown using RT-PCR Primer: ELP6 (m)-PR: sc-154341-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.